

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal652dmr

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page for STN Seminar Schedule - N. America  
NEWS 2 NOV 21 CAS patent coverage to include exemplified prophetic  
substances identified in English-, French-, German-,  
and Japanese-language basic patents from 2004-present  
NEWS 3 NOV 26 MARPAT enhanced with FSORT command  
NEWS 4 NOV 26 CHEMSAFE now available on STN Easy  
NEWS 5 NOV 26 Two new SET commands increase convenience of STN  
searching  
NEWS 6 DEC 01 ChemPort single article sales feature unavailable  
NEWS 7 DEC 12 GBFULL now offers single source for full-text  
coverage of complete UK patent families  
NEWS 8 DEC 17 Fifty-one pharmaceutical ingredients added to FS  
NEWS 9 JAN 06 The retention policy for unread STNmail messages  
will change in 2009 for STN-Columbus and STN-Tokyo  
NEWS 10 JAN 07 WPIIDS, WPINDEXT, and WPIX enhanced Japanese Patent  
Classification Data  
NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,  
AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS LOGIN Welcome Banner and News Items  
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 00:49:28 ON 01 FEB 2009

=> index bioscience medicine

FILE 'DRUGMONOG' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.22	0.22

FULL ESTIMATED COST

INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, ANTE, AQUALINE,  
AQUASCI, BIORNG, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CAB, CAPLUS,  
CEABA-VTB, CIN, CONFSCI, CROPB, CROPU, DDPB, DDFU, DGENE, DISSABS, DRUGB,

71 FILES IN THE FILE LIST IN STNINDEX

Enter SET DETAIL ON to see search term postings or to view  
search error messages that display as 0\* with SET DETAIL OFF.

-> s (restric?(3a)endonucleas?) or (restric?(3a)enzym?) or  
(restric?(3a)modif?(5a) (enzym? or endonucleas? or system?))

21 FILE ADISCTI  
3 FILE ADISINSIGHT  
7 FILE ADISNEWS  
3845 FILE AGRICOLA  
104 FILE ANABSTR  
42 FILE ANTE  
40 FILE AQUALINE  
1189 FILE AQUASCI  
3479 FILE BIOENG

9 FILES SEARCHED...  
31117 FILE BIOSIS  
10212 FILE BIOTECHABS  
10212 FILE BIOTECHDS  
17093 FILE BIOTECHNO

13 FILES SEARCHED...  
10248 FILE CABA  
40802 FILE CAPLUS  
731 FILE CEABA-VTB  
77 FILE CIN  
338 FILE CONFSCI  
3 FILE CROPB  
124 FILE CROPU  
19 FILE DDFB  
133 FILE DDFU  
43989 FILE DGENE

23 FILES SEARCHED...  
2088 FILE DISSABS  
19 FILE DRUGB  
425 FILE DRUGU  
88 FILE EMBAL  
21832 FILE EMBASE  
9157 FILE ESBIODASE

30 FILES SEARCHED...  
363 FILE PROSTI  
1154 FILE PSTA  
2282317 FILE GENBANK  
53 FILE HEALSAFE  
7069 FILE IFIPAT  
11 FILE IMSDRUGNEWS

39 FILES SEARCHED...  
9 FILE IMSRESEARCH  
20 FILE KOSMET  
17321 FILE LIFESCI  
40206 FILE MEDLINE  
199 FILE NTIS  
348 FILE OCEAN  
11890 FILE PASCAL

47 FILES SEARCHED...  
164 FILE PCTGEN  
1 FILE PHAR  
1 FILE PHARMAML  
66 FILE PHIN  
640 FILE PROMT

```

      1  FILE PROUSDDR
      3  FILE RDISCLOSURE
20911  FILE SCISEARCH
10529  FILE TOXCENTER
58 FILES SEARCHED...
14486  FILE USGENE
72194  FILE USPATFULL
      29  FILE USPATOLD
11872  FILE USPAT2
      1  FILE VETB
63 FILES SEARCHED...
      144  FILE VETU
      62  FILE WATER
9109   FILE WPIDS
      62  FILE WPIFV
9109   FILE WPINDEX
      37  FILE IPA
      4  FILE NAPRALERT
488    FILE NLDB

```

64 FILES HAVE ONE OR MORE ANSWERS, 71 FILES SEARCHED IN STINDEX

L1 QUE (RESTRIC?(3A) ENDONUCLEAS?) OR (RESTRIC?(3A) ENZYM?) OR (RESTRIC?(3A) MODIF?(5A) (ENZYM? OR ENDONUCLEAS? OR SYSTEM?))

=> d rank

```

F1 2282317 GENBANK
F2 72194   USPATFULL
F3 43989   DGENE
F4 40802   CAPLUS
F5 40206   MEDLINE
F6 31117   BIOSIS
F7 21832   EMBASE
F8 20911   SCISEARCH
F9 17321   LIFESCI
F10 17093  BIOTECHNO
F11 14486  USGENE
F12 11890  PASCAL
F13 11872  USPAT2
F14 10529  TOXCENTER
F15 10248  CABA
F16 10212  BIOTECHABS
F17 10212  BIOTECHDS
F18 9157   ESBIOBASE
F19 9109   WPIDS
F20 9109   WPINDEX
F21 7069   IPIPAT
F22 3845   AGRICOLA
F23 3479   BIOENG
F24 2088   DISSABS
F25 1189   AQUASCI
F26 1154   FSTA
F27 731    CEABA-VTB
F28 640    PROMT
F29 488    NLDB
F30 425    DRUGU
F31 363    FROSTI
F32 348    OCEAN
F33 338    CONFSCI
F34 199    NTIS
F35 164    PCTGEN
F36 144    VETU

```

F37	133	DDPU
F38	124	CROPU
F39	104	ANABSTR
F40	88	EMBAL
F41	77	CIN
F42	66	PHIN
F43	62	WATER
F44	62	WPIFV
F45	53	HEALSAFE
F46	42	ANTE
F47	40	AQUALINE
F48	37	IPA
F49	29	USPATOLD
F50	21	ADISCTI
F51	20	KOSMET
F52	19	DDFB
F53	19	DRUGB
F54	11	IMSDRUGNEWS
F55	9	IMSRESEARCH
F56	7	ADISNEWS
F57	4	NAPRALERT
F58	3	ADISINSIGHT
F59	3	CROPB
F60	3	RDISCLOSURE
F61	1	PHAR
F62	1	PHARMAML
F63	1	PROUSDDR
F64	1	VETB

-> file f2, f4-f17

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	12.24	12.46

FILE 'USPATFULL' ENTERED AT 01:00:54 ON 01 FEB 2009  
CA INDEXING COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'CAPLOS' ENTERED AT 01:00:54 ON 01 FEB 2009  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 01:00:54 ON 01 FEB 2009

FILE 'BIOSIS' ENTERED AT 01:00:54 ON 01 FEB 2009  
Copyright (c) 2009 The Thomson Corporation

FILE 'EMBASE' ENTERED AT 01:00:54 ON 01 FEB 2009  
Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'SCISEARCH' ENTERED AT 01:00:54 ON 01 FEB 2009  
Copyright (c) 2009 The Thomson Corporation

FILE 'LIFESCI' ENTERED AT 01:00:54 ON 01 FEB 2009  
COPYRIGHT (C) 2009 Cambridge Scientific Abstracts (CSA)

FILE 'BIOTECHNO' ENTERED AT 01:00:54 ON 01 FEB 2009  
COPYRIGHT (C) 2009 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'USGENE' ENTERED AT 01:00:54 ON 01 FEB 2009

COPYRIGHT (C) 2009 SEQUENCEBASE CORP

FILE 'PASCAL' ENTERED AT 01:00:54 ON 01 FEB 2009

Any reproduction or dissemination in part or in full,  
by means of any process and on any support whatsoever  
is prohibited without the prior written agreement of INIST-CNRS.  
COPYRIGHT (C) 2009 INIST-CNRS. All rights reserved.

FILE 'USPAT2' ENTERED AT 01:00:54 ON 01 FEB 2009

CA INDEXING COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'TOXCENTER' ENTERED AT 01:00:54 ON 01 FEB 2009

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'CABA' ENTERED AT 01:00:54 ON 01 FEB 2009

COPYRIGHT (C) 2009 CAB INTERNATIONAL (CABI)

FILE 'BIOTECHABS' ACCESS NOT AUTHORIZED

FILE 'BIOTECHDS' ENTERED AT 01:00:54 ON 01 FEB 2009

COPYRIGHT (C) 2009 THOMSON REUTERS

=> s (restric?(3a)endonucleas?) or (restric?(3a)enzym?) or  
(restric?(3a)modif?(5a)(enzym? or endonucleas? or system?))

7 FILES SEARCHED...

10 FILES SEARCHED...

L2 330713 (RESTRIC?(3A) ENDONUCLEAS?) OR (RESTRIC?(3A) ENZYM?) OR (RESTRIC  
?(3A) MODIF?(5A) ENZYM? OR ENDONUCLEAS? OR SYSTEM?))

=> s l2(s)(specifi? or recog?)(s)(sequenc? or dna?)

5 FILES SEARCHED...

9 FILES SEARCHED...

L3 79162 L2(S)(SPECIFI? OR RECOG?)(S)(SEQUENC? OR DNA?)

=> s l3 and (hybrid? or recombinat? or truncat? or transpos?)

L4 49041 L3 AND (HYBRID? OR RECOMBINAT? OR TRUNCAT? OR TRANSPOS?)

=> s l3(s)((two(3a)recognit?(3a)site?) or hsd?)

L5 990 L3(S)((TWO(3A) RECOGNIT?(3A) SITE?) OR HSDS?)

=>

=> s l5 (s)(hybrid? or recombin? or trunca? or or exchang? or transpos? or alter?)  
MISSING TERM 'OR OR'

The search profile that was entered contains a logical  
operator followed immediately by another operator.

=> s l5 (s)(hybrid? or recombin? or trunca? or exchang? or transpos? or alter?)

5 FILES SEARCHED...

11 FILES SEARCHED...

L6 344 L5 (S)(HYBRID? OR RECOMBIN? OR TRUNCA? OR EXCHANG? OR TRANSPOS?  
OR ALTER?)

=> dup rem l6

DUPLICATE IS NOT AVAILABLE IN 'USGENE'.

ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE  
PROCESSING COMPLETED FOR L6

L7 280 DUP REM L6 (64 DUPLICATES REMOVED)

=> s l7(s)half?

PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH  
FIELD CODE - 'AND' OPERATOR ASSUMED 'L&2(S)HALF?'  
PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH

FIELD CODE - 'AND' OPERATOR ASSUMED 'L86(S)HALF?'  
 PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH  
 FIELD CODE - 'AND' OPERATOR ASSUMED 'L88(S)HALF?'  
 PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH  
 FIELD CODE - 'AND' OPERATOR ASSUMED 'L100(S)HALF?'  
 PROXIMITY OPERATOR LEVEL NOT CONSISTENT WITH  
 FIELD CODE - 'AND' OPERATOR ASSUMED 'L102(S)HALF?'  
 L8 14 L7(S) HALF?

=> d ti l8 1-14

L8 ANSWER 1 OF 14 USPATFULL on STN  
 TI Small interfering RNA libraries and methods of synthesis and use

L8 ANSWER 2 OF 14 USPATFULL on STN  
 TI Nucleic acid and amino acid sequences relating to Enterobacter cloacae for diagnostics and therapeutics

L8 ANSWER 3 OF 14 USPATFULL on STN  
 TI Differential enzymatic fragmentation by whole genome amplification

L8 ANSWER 4 OF 14 USPATFULL on STN  
 TI Differential enzymatic fragmentation

L8 ANSWER 5 OF 14 USPATFULL on STN  
 TI Methods for quantitative determination of methylation density in a DNA locus

L8 ANSWER 6 OF 14 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of colon cancer

L8 ANSWER 7 OF 14 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of pancreatic cancer

L8 ANSWER 8 OF 14 USPATFULL on STN  
 TI Nuclease

L8 ANSWER 9 OF 14 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of colon cancer

L8 ANSWER 10 OF 14 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of ovarian cancer

L8 ANSWER 11 OF 14 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of colon cancer

L8 ANSWER 12 OF 14 LIFESCI COPYRIGHT 2009 CSA on STN  
 TI DNA Cleavage by Type III Restriction-modification Enzyme Eco P15I is Independent of Spacer Distance between Two Head to Head Oriented Recognition Sites

L8 ANSWER 13 OF 14 LIFESCI COPYRIGHT 2009 CSA on STN  
 TI Generation of new DNA binding specificity by truncation of the type IC EcoDXXI hsdS gene

L8 ANSWER 14 OF 14 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI Dendrimer based multifunctional composition for treating cancer and cardiovascular disease, comprises a dendrimer complex having dendrimers comprising different agents e.g. therapeutic and biological monitoring agents;  
 useful for inflammatory disease and pathogen disease gene therapy,

diagnosis and drug target screening

-> d 18 ibib abs 8 12 13

L8 ANSWER 8 OF 14 USPATFULL on STN

ACCESSION NUMBER: 2003:58047 USPATFULL  
 TITLE: Nuclease  
 INVENTOR(S): Janulaitis, Arvydas, Vilnius, LITHUANIA  
 Rimseleiene, Renata, Vilnius, LITHUANIA  
 Lubys, Arvydas, Vilnius, LITHUANIA  
 PATENT ASSIGNEE(S): Fermentas AB (non-U.S. corporation)

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 20030040614	A1	20030227
	US 6893854	B2	20050517
APPLICATION INFO.:	US 2001-906768	A1	20010718 (9)

	NUMBER	DATE
PRIORITY INFORMATION:	GB 2000-19744	20000810
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	APPLICATION	
LEGAL REPRESENTATIVE:	PILLSBURY WINTHROP, LLP, P.O. BOX 10500, MCLEAN, VA, 22102	
NUMBER OF CLAIMS:	17	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	6 Drawing Page(s)	
LINE COUNT:	1029	

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB A process for producing a polynucleotide encoding a restriction endonuclease with an altered specificity, which process comprises:

(a) mutagenising a polynucleotide encoding a restriction endonuclease with specificity for a recognition sequence so as to produce one or more mutated polynucleotides; and

(b) isolating therefrom a polynucleotide encoding a mutated restriction endonuclease with specificity for an altered recognition sequence by selecting a polynucleotide which expresses a restriction endonuclease with methylase specificity for the altered recognition sequence.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L8 ANSWER 12 OF 14 LIFESCI COPYRIGHT 2009 CSA on STN

ACCESSION NUMBER: 2001:107290 LIFESCI  
 TITLE: DNA Cleavage by Type III Restriction-modification Enzyme Eco P15I is Independent of Spacer Distance between Two Head to Head Oriented Recognition Sites  
 AUTHOR: Muecke, M.; Reich, S.; Moencke-Buchner, E.; Reuter, M.; Krueger, D.H.\*  
 CORPORATE SOURCE: Institut fuer Virologie, Medizinische Fakultät (Charité), der Humboldt-Universität zu Berlin, D-10098, Berlin, Germany  
 SOURCE: Journal of Molecular Biology [J. Mol. Biol.], (20010928) vol. 312, no. 4, pp. 687-698.  
 ISSN: 0022-2836.  
 DOCUMENT TYPE: Journal  
 FILE SEGMENT: N; J  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB The type III restriction-modification enzyme EcoP15I requires the interaction of two unmethylated, inversely oriented recognition sites 5'-CAGCAG in head to head configuration to allow an efficient DNA cleavage. It has been hypothesized that two convergent DNA-translocating enzyme-substrate complexes interact to form the active cleavage complex and that translocation is driven by ATP hydrolysis. Using a half-automated, fluorescence-based detection method, we investigated how the distance between two inversely oriented recognition sites affects DNA cleavage efficiency. We determined that EcoP15I cleaves DNA efficiently even for two adjacent head to head or tail to tail oriented target sites. Hence, DNA translocation appears not to be required for initiating DNA cleavage in these cases. Furthermore, we report here that EcoP15I is able to cleave single-site substrates. When we analyzed the interaction of EcoP15I with DNA substrates containing adjacent target sites in the presence of non-hydrolyzable ATP analogues, we found that cleavage depended on the hydrolysis of ATP. Moreover, we show that cleavage occurs at only one of the two possible cleavage positions of an interacting pair of target sequences. When EcoP15I bound to a DNA substrate containing one recognition site in the absence of ATP, we observed a 36 nucleotide DNaseI-footprint that is asymmetric on both strands. All of our footprinting experiments showed that the enzyme did not cover the region around the cleavage site. Analyzing a DNA fragment with two head to head oriented recognition sites, EcoP15I protected 27-33 nucleotides around the recognition sequence, including an additional region of 26 bp between both cleavage sites. For all DNA substrates examined, the presence of ATP caused altered footprinting patterns. We assume that the altered patterns are most likely due to a conformational change of the enzyme. Overall, our data further refine the tracking-collision model for type III restriction enzymes. Copyright 2001 Academic Press

L8 ANSWER 13 OF 14 LIFESCI COPYRIGHT 2009 CSA on STN

ACCESSION NUMBER: 97:330 LIFESCI  
 TITLE: Generation of new DNA binding specificity by truncation of the type IC EcoDXXI hsdS gene  
 AUTHOR: MacWilliams, M.P.; Bickle, T.A.\*  
 CORPORATE SOURCE: Dep. Microbiol., Biozentrum, Basel Univ., Klingelbergstrasse 70, CH-4056 Basel, Switzerland  
 SOURCE: EMBO J., (1996) vol. 15, no. 17, pp. 4775-4783. ISSN: 0261-4189.  
 DOCUMENT TYPE: Journal  
 FILE SEGMENT: G; J  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB The hsdS subunit of a type IC restriction-modification enzyme is responsible for the enzyme's DNA binding specificity. Type I recognition sites are characterized by two defined half-sites separated by a non-specific spacer of defined length. The hsdS subunit contains two independent DNA binding domains, each targeted towards one DNA half-site. We have shown previously that the 5' half of hsdS can code for a functional substitute of the full-length hsdS. Here we demonstrate that the 3' half of the gene, when fused to the appropriate transcriptional and translational start signals, also codes for a peptide which imparts DNA binding specificity to the enzyme. About half the natural hsdS size, the mutant peptide contains a single DNA recognition domain flanked by one copy of each internal repeat found in the



full-length hsdS. Deletion of either repeat sequence results in loss of activity. Like the 5' hsdS mutant, the 3' mutant recognizes an interrupted palindrome, GAAYN sub(5)RTTC, suggesting that two truncated subunits participate in DNA recognition. Coexpression of the 5' hsdS mutant and the 3' hsdS mutant along with hsdM regenerates the wild-type methylation specificity. Thus, there is a free assortment of subunits in the cell.

```
-> s l2(s)((two(3a)recog?(3a)site?) or hsdS? or (half?(3a)site?))
L9      1657 L2(S)((TWO(3A) RECOG?(3A) SITE?) OR HSDS? OR (HALF?(3A) SITE?))

-> s l9(s)(hybrid? or recomb? or trunc? or exch? or transpo? or alter?)
  10 FILES SEARCHED...
L10      500 L9(S)(HYBRID? OR RECOMB? OR TRUNC? OR EXCH? OR TRANSP? OR
      ALTER?)

-> s l10 and ((modif? or alter? or hybrid?) (3a)(dna? or sequen? or specific? or
(recogn?(3a)site?)))
  1 FILES SEARCHED...
  2 FILES SEARCHED...
  8 FILES SEARCHED...
  9 FILES SEARCHED...
 10 FILES SEARCHED...
 11 FILES SEARCHED...
L11      359 L10 AND ((MODIF? OR ALTER? OR HYBRID?) (3A) (DNA? OR SEQUEN? OR
      SPECIFIC? OR (RECOGN?(3A) SITE?)))

-> dup rem l11
DUPLICATE IS NOT AVAILABLE IN 'USGENE'.
ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE
PROCESSING COMPLETED FOR L11
L12      314 DUP REM L11 (45 DUPLICATES REMOVED)

-> d ti l12 1-314

L12 ANSWER 1 OF 314  USPATFULL on STN
TI   HOMOLOGOUS RECOMBINATION IN PLANTS

L12 ANSWER 2 OF 314  USPATFULL on STN
TI   Methods and Compositions Involving Polymeric Immunoglobulin Fusion
      Proteins

L12 ANSWER 3 OF 314  USPATFULL on STN
TI   Rabies Virus Vector Systems and Compositions and Methods Thereof

L12 ANSWER 4 OF 314  USPATFULL on STN
TI   Polypeptides from Non-Typeable Haemophilus Influenzae

L12 ANSWER 5 OF 314  USPATFULL on STN
TI   Analysis of methylation using selective adaptor ligation

L12 ANSWER 6 OF 314  USPATFULL on STN
TI   Methods for Identification of Merle Gene

L12 ANSWER 7 OF 314  USPATFULL on STN
TI   Compositions and Methods for Genetic Manipulation and Monitoring of Cell
      Lines

L12 ANSWER 8 OF 314  USPATFULL on STN
```

TI Selection and Enrichment of Proteins Using in vitro Compartmentalization

L12 ANSWER 9 OF 314 USPATFULL on STN

TI OLIGONUCLEOTIDE LINKERS COMPRISING A VARIABLE COHESIVE PORTION AND METHOD FOR THE PREPARATION OF POLYNUCLEOTIDE LIBRARIES BY USING SAID LINKERS

L12 ANSWER 10 OF 314 USPATFULL on STN

TI Attenuated parainfluenza virus (PIV) vaccines

L12 ANSWER 11 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN

TI Expressing a sequence of interest in a plant cell, including a plant, plant part, or plant cell culture comprises providing a cell with a DNA sequence and causing expression of the sequence of interest; recombinant protein produced by vector mediated gene expression in host cell, useful in construction of transgenic plant

L12 ANSWER 12 OF 314 USPATFULL on STN DUPLICATE 1

TI METHODS FOR IDENTIFICATION OF ALPORT SYNDROME

L12 ANSWER 13 OF 314 USPATFULL on STN DUPLICATE 2

TI Nucleic acid and amino acid sequences relating to streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 14 OF 314 USPATFULL on STN DUPLICATE 3

TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 15 OF 314 USPATFULL on STN DUPLICATE 4

TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 16 OF 314 USPATFULL on STN

TI NON-REDUCING SACCHARIDE-FORMING ENZYME, TREHALOSE-RELEASING ENZYME, AND PROCESS FOR PRODUCING SACCHARIDES USING THE ENZYMES

L12 ANSWER 17 OF 314 USPATFULL on STN

TI Group b streptococcus antigens

L12 ANSWER 18 OF 314 USPATFULL on STN

TI Artificial plant minichromosomes

L12 ANSWER 19 OF 314 USPATFULL on STN

TI Methods for assembly of high fidelity synthetic polynucleotides

L12 ANSWER 20 OF 314 USPATFULL on STN

TI Methods and Means for Regulating Gene Expression

L12 ANSWER 21 OF 314 USPATFULL on STN

TI Directed enrichment of genomic DNA for high-throughput sequencing

L12 ANSWER 22 OF 314 USPATFULL on STN

TI NON-REDUCING SACCHARIDE-FORMING ENZYME, TREHALOSE-RELEASING ENZYME, AND PROCESS FOR PRODUCING SACCHARIDES USING THE ENZYMES

L12 ANSWER 23 OF 314 USPATFULL on STN

TI Small interfering RNA libraries and methods of synthesis and use

L12 ANSWER 24 OF 314 USPATFULL on STN

TI Novel modular type II restriction endonuclease, cspci, and the use of modular endonucleases for generating endonucleases with new

specificities

- L12 ANSWER 25 OF 314 USPATFULL on STN  
TI Targeted integration and expression of exogenous nucleic acid sequences
- L12 ANSWER 26 OF 314 USPATFULL on STN  
TI Attenuated human-bovine chimeric parainfluenza virus (PIV) vaccines
- L12 ANSWER 27 OF 314 USPATFULL on STN  
TI Methods for assembly of high fidelity synthetic polynucleotides
- L12 ANSWER 28 OF 314 USPATFULL on STN  
TI BIOINFORMATICALLY DETECTABLE GROUP OF NOVEL VACCINIA REGULATORY GENES AND USES THEREOF
- L12 ANSWER 29 OF 314 USPATFULL on STN  
TI Process for chromosomal expression of foreign genes in the hsdM region of a methylotrophic microbial host cell
- L12 ANSWER 30 OF 314 USPATFULL on STN  
TI Methods for genotyping
- L12 ANSWER 31 OF 314 USPATFULL on STN  
TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics
- L12 ANSWER 32 OF 314 USPATFULL on STN  
TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics
- L12 ANSWER 33 OF 314 USPATFULL on STN  
TI Cell line and methods for determining viral titer
- L12 ANSWER 34 OF 314 USPATFULL on STN  
TI Insertion sequence-free bacteria
- L12 ANSWER 35 OF 314 USPATFULL on STN  
TI Bacteria with reduced genome
- L12 ANSWER 36 OF 314 USPATFULL on STN  
TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics
- L12 ANSWER 37 OF 314 USPATFULL on STN  
TI Evolution of whole cells and organisms by recursive sequence recombination
- L12 ANSWER 38 OF 314 USPATFULL on STN  
TI Methods for producing polypeptide-tagged collections and capture systems containing the tagged polypeptides
- L12 ANSWER 39 OF 314 USPATFULL on STN  
TI Isoprenoid biosynthesis
- L12 ANSWER 40 OF 314 USPATFULL on STN  
TI Methods for monitoring multiple gene expression
- L12 ANSWER 41 OF 314 USPATFULL on STN  
TI Nucleic acid and amino acid sequences relating to streptococcus pneumoniae for diagnostics and therapeutics
- L12 ANSWER 42 OF 314 USPATFULL on STN

TI Nucleic acid and amino acid sequences relating to streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 43 OF 314 USPATFULL on STN

TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 44 OF 314 USPATFULL on STN

TI Plasmids and methods for construction of non-redundant, indexed, saturation, gene-disruption plant and animal libraries

L12 ANSWER 45 OF 314 USPATFULL on STN

TI Attenuated human-bovine chimeric parainfluenza virus(PIV) vaccines

L12 ANSWER 46 OF 314 USPATFULL on STN

TI Non-reducing saccharide-forming enzyme, trehalose-releasing enzyme, and process for producing saccharides using the enzymes

L12 ANSWER 47 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN

TI Juxtaposing sequence tags comprises digesting a DNA adaptor to cleave the target DNA insert to create two sequence tags comprising terminal sequences of the target DNA insert that are attached to the plasmid vector;  
juxtaposing sequence tag via DNA adaptor digestion for disease diagnosis and genomics

L12 ANSWER 48 OF 314 USPATFULL on STN DUPLICATE 5

TI Anthranilate synthase gene and method for increasing tryptophan production

L12 ANSWER 49 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 6

TI Polypeptide mutagenesis method using transposons containing restriction enzyme recognition sites toward each termini of the encoding DNA target

L12 ANSWER 50 OF 314 USPATFULL on STN

TI Analysis of methylation using nucleic acid arrays

L12 ANSWER 51 OF 314 USPATFULL on STN

TI Accessible polynucleotide libraries and methods of use thereof

L12 ANSWER 52 OF 314 USPATFULL on STN

TI Gram positive bacterial mutants and methods of generating and using such mutants

L12 ANSWER 53 OF 314 USPATFULL on STN

TI Bacteria with reduced genome

L12 ANSWER 54 OF 314 USPATFULL on STN

TI NOVEL POLYNUCLEOTIDES ENCODING USEFUL POLYPEPTIDES IN CORYNEBACTERIUM GLUTAMICUM SSP. LACTOFERMENTUM

L12 ANSWER 55 OF 314 USPATFULL on STN

TI NOVEL POLYNUCLEOTIDES ENCODING USEFUL POLYPEPTIDES IN CORYNEBACTERIUM GLUTAMICUM SSP. LACTOFERMENTUM

L12 ANSWER 56 OF 314 USPATFULL on STN

TI Strain belonging to the genus streptomyces and being capable of producing nemadictin and process for producing nemadictin using the strain

L12 ANSWER 57 OF 314 USPATFULL on STN

TI Methods and compounds for raising antibodies and for screening antibody

repertoires

- L12 ANSWER 58 OF 314 USPATFULL on STN  
TI Insertion Sequence-Free Bacteria
- L12 ANSWER 59 OF 314 USPATFULL on STN  
TI Methods for assembly of high fidelity synthetic polynucleotides
- L12 ANSWER 60 OF 314 USPATFULL on STN  
TI Method for the manufacture of nucleic acid molecules
- L12 ANSWER 61 OF 314 USPATFULL on STN  
TI Polypeptides and polynucleotides from coagulase-negative staphylococci
- L12 ANSWER 62 OF 314 USPATFULL on STN  
TI Protein having PDZ domain sequence
- L12 ANSWER 63 OF 314 USPATFULL on STN  
TI Methods and compositions for elucidating protein expression profiles in cells
- L12 ANSWER 64 OF 314 USPATFULL on STN  
TI Attenuated human-bovine chimeric parainfluenza virus (PIV) vaccines
- L12 ANSWER 65 OF 314 USPATFULL on STN  
TI Hybrid and single chain meganucleases and use thereof
- L12 ANSWER 66 OF 314 USPATFULL on STN  
TI Methods and systems for in silico experimental design and for providing a biotechnology product to a customer
- L12 ANSWER 67 OF 314 USPATFULL on STN  
TI Effect of treatment with 4,5-dihydroxy-2-cyclopenten-1-one (dhcp) on gene expression and quorum-sensing in bacteria
- L12 ANSWER 68 OF 314 USPATFULL on STN  
TI Peptides for metal ion affinity chromatography
- L12 ANSWER 69 OF 314 USPATFULL on STN  
TI Analysis of mixtures of nucleic acid fragments
- L12 ANSWER 70 OF 314 USPATFULL on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof
- L12 ANSWER 71 OF 314 USPATFULL on STN  
TI Methods of producing mutant polynucleotides
- L12 ANSWER 72 OF 314 USPATFULL on STN  
TI Sequence specific recombinase-based methods for producing intron containing vectors and compositions for use in practicing the same
- L12 ANSWER 73 OF 314 USPATFULL on STN  
TI Isoprenoid biosynthesis
- L12 ANSWER 74 OF 314 USPATFULL on STN  
TI Nucleic acid and amino acid sequences relating to *Enterobacter cloacae* for diagnostics and therapeutics
- L12 ANSWER 75 OF 314 USPATFULL on STN  
TI Protein having PDZ domain sequence

L12 ANSWER 76 OF 314 USPATFULL on STN  
 TI Cystic fibrosis gene

L12 ANSWER 77 OF 314 USPAT2 on STN  
 TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 78 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI New isolated nucleotide sequence of pgm gene of a Brucella bacterium modified by a partial deletion of the sequence coding for phosphoglucomutase, useful for producing vaccines for treating brucellosis;  
 recombinant vaccine preparation via attenuated bacterium strain for use in disease therapy

L12 ANSWER 79 OF 314 LIFESCI COPYRIGHT 2009 CSA on STN  
 TI Site-specific labeling of supercoiled DNA

L12 ANSWER 80 OF 314 USPATFULL on STN DUPLICATE 7  
 TI Differential enzymatic fragmentation by whole genome amplification

L12 ANSWER 81 OF 314 USPATFULL on STN  
 TI 2', 5'-oligoadenylate phosphodiesterase

L12 ANSWER 82 OF 314 USPATFULL on STN  
 TI Novel oligonucleotide arrays and their use for sorting, isolating, sequencing, and manipulating nucleic acids

L12 ANSWER 83 OF 314 USPATFULL on STN  
 TI Transposon

L12 ANSWER 84 OF 314 USPATFULL on STN  
 TI Generation and application of standardized universal libraries

L12 ANSWER 85 OF 314 USPATFULL on STN  
 TI Nucleic acid molecules containing recombination sites and methods of using the same

L12 ANSWER 86 OF 314 USPATFULL on STN  
 TI Targeted chromosomal mutagenesis using zinc finger nucleases

L12 ANSWER 87 OF 314 USPATFULL on STN  
 TI Analysis of methylation status using nucleic acid arrays

L12 ANSWER 88 OF 314 USPATFULL on STN  
 TI Differential enzymatic fragmentation

L12 ANSWER 89 OF 314 USPATFULL on STN  
 TI Analysis of methylation status using oligonucleotide arrays

L12 ANSWER 90 OF 314 USPATFULL on STN  
 TI Methods for quantitative determination of methylation density in a DNA locus

L12 ANSWER 91 OF 314 USPATFULL on STN  
 TI Method for comprehensive identification of cell lineage specific genes

L12 ANSWER 92 OF 314 USPATFULL on STN  
 TI Tryparedoxin, expression plasmid, process of preparation, method of use, test kit and pharmaceutical composition

L12 ANSWER 93 OF 314 USPATFULL on STN

TI Nucleic acid and amino acid sequences relating to streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 94 OF 314 USPATFULL on STN

TI Nucleotide sequence of the haemophilus influenzae Rd genome, fragments thereof, and uses thereof

L12 ANSWER 95 OF 314 USPATFULL on STN

TI Methods and compositions for detecting promoter activity and expressing fusion proteins

L12 ANSWER 96 OF 314 USPATFULL on STN

TI Cloning vectors and method for molecular cloning

L12 ANSWER 97 OF 314 USPATFULL on STN

TI Methods for insertion of nucleic acids into circular vectors

L12 ANSWER 98 OF 314 USPATFULL on STN

TI Systems for capture and analysis of biological particles and methods using the systems

L12 ANSWER 99 OF 314 USPATFULL on STN

TI Mapping genomic rearrangements

L12 ANSWER 100 OF 314 USPATFULL on STN

TI Concurrent enzymatic polynucleotide synthesis and detectable signal generation

L12 ANSWER 101 OF 314 USPATFULL on STN

TI Methods of diagnosing and treating hepatic cell proliferative disorders

L12 ANSWER 102 OF 314 USPATFULL on STN

TI Analysis of methylation status using oligonucleotide arrays

L12 ANSWER 103 OF 314 USPATFULL on STN

TI Plant promoter and method for gene expression using said promoter

L12 ANSWER 104 OF 314 USPATFULL on STN

TI Cystic fibrosis gene

L12 ANSWER 105 OF 314 USP2 on STN

TI Nucleic acid amplification method

L12 ANSWER 106 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN

TI New pure Type IIG restriction endonuclease obtainable from Citrobacter species or from Escherichia coli, useful for generating restriction endonucleases with new specificities;  
for use in genetic engineering

L12 ANSWER 107 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN

TI Identifying a conditional essential gene of an organism, useful for manufacturing a medicament for vaccinating a human or animal, comprises providing a library of transposon mutants of the organism;  
involving vector-mediated gene transfer and expression in host cell  
for use in therapy and recombinant vaccine preparation

L12 ANSWER 108 OF 314 USPATFULL on STN DUPLICATE 8

TI Group b streptococcus antigens and corresponding dna fragments

L12 ANSWER 109 OF 314 USPATFULL on STN DUPLICATE 9

TI Novel polynucleotides encoding useful polypeptides in corynebacterium glutamicum SSP. lactofermentum

L12 ANSWER 110 OF 314 USPATFULL on STN DUPLICATE 10  
 TI Use of multiple recombination sites with unique specificity in recombinational cloning

L12 ANSWER 111 OF 314 USPATFULL on STN DUPLICATE 11  
 TI NUCLEOTIDE SEQUENCE OF THE HAEMOPHILUS INFLUENZAE RD GENOME, FRAGMENTS THEREOF, AND USES THEREOF

L12 ANSWER 112 OF 314 USPATFULL on STN DUPLICATE 12  
 TI Proteins and polypeptides from coagulase-negative staphylococci

L12 ANSWER 113 OF 314 USPATFULL on STN DUPLICATE 13  
 TI Non-mevalonate isoprenoid pathway

L12 ANSWER 114 OF 314 USPATFULL on STN  
 TI Use of site specific recombination to prepare molecular markers

L12 ANSWER 115 OF 314 USPATFULL on STN  
 TI Methods and compositions relating to 5'-chimeric ribonucleic acids

L12 ANSWER 116 OF 314 USPATFULL on STN  
 TI Viral vectors containing recombination sites

L12 ANSWER 117 OF 314 USPATFULL on STN  
 TI Methods for producing polypeptide-tagged collections and capture systems containing the tagged polypeptides

L12 ANSWER 118 OF 314 USPATFULL on STN  
 TI Subscription based systems, methods and components for providing genomic and proteomic products and services

L12 ANSWER 119 OF 314 USPATFULL on STN  
 TI Method for high throughput elucidation of transcriptional profiles and genome annotation

L12 ANSWER 120 OF 314 USPATFULL on STN  
 TI Antigens of group b streptococcus and corresponding dna fragments

L12 ANSWER 121 OF 314 USPATFULL on STN  
 TI Oligonucleotide linkers comprising a variable cohesive portion and method for the preparation of polynucleotide libraries by using said linkers

L12 ANSWER 122 OF 314 USPATFULL on STN  
 TI Methods for treating or preventing infections from coagulase-negative staphylococci

L12 ANSWER 123 OF 314 USPATFULL on STN  
 TI Nucleotide sequences of moraxella catarrhalis genome

L12 ANSWER 124 OF 314 USPATFULL on STN  
 TI Human complement C3-binding protein from streptococcus pneumoniae

L12 ANSWER 125 OF 314 USPATFULL on STN  
 TI Novel recombinant xylanases derived from anaerobic fungi, and the relevant sequences, expression vectors and hosts

L12 ANSWER 126 OF 314 USPATFULL on STN  
 TI Use of collections of binding sites for sample profiling and other applications



L12 ANSWER 127 OF 314 USPTAFULL on STN  
 TI Antibodies to polypeptides from coagulase-negative staphylococci

L12 ANSWER 128 OF 314 USPTAFULL on STN  
 TI Nucleotide sequence of the haemophilus influenza Rd genome, fragments thereof, and uses thereof

L12 ANSWER 129 OF 314 USPTAFULL on STN  
 TI Modification of plant cell wall component and method of regulating development differentiation

L12 ANSWER 130 OF 314 USPTAFULL on STN  
 TI Method for the identification of essential and conditional essential genes

L12 ANSWER 131 OF 314 USPTAFULL on STN  
 TI Molecular diagnosis of bacteremia

L12 ANSWER 132 OF 314 USPTAFULL on STN  
 TI Hybrid and single chain meganucleases and use thereof

L12 ANSWER 133 OF 314 USPTAFULL on STN  
 TI DNA sequences, recombinant DNA molecules and processes for producing human interferon-like polypeptides

L12 ANSWER 134 OF 314 USPTAFULL on STN  
 TI Methods for insertion of nucleic acids into circular vectors

L12 ANSWER 135 OF 314 USPTAFULL on STN  
 TI Nucleic acid and amino acid sequences relating to Streptococcus pneumoniae for diagnostics and therapeutics

L12 ANSWER 136 OF 314 USPTAFULL on STN  
 TI Cystic fibrosis gene

L12 ANSWER 137 OF 314 USPTAFULL on STN  
 TI Methods of diagnosing and treating hepatic cell proliferative disorders

L12 ANSWER 138 OF 314 USPTAFULL on STN  
 TI Human complement C3-degrading protein from Streptococcus pneumoniae

L12 ANSWER 139 OF 314 USPAT2 on STN  
 TI NCC2705--the genome of a bifidobacterium

L12 ANSWER 140 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI Preparing small interfering RNA library for treating e.g. cancer, by producing random oligoDNAs that can be cloned into vectors containing site-specific recombinase sites for generating inverted repeats of the sequence in host cells;  
     for use in cancer prevention, gene therapy, RNA interference and functional genomics

L12 ANSWER 141 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI Identifying sequences that express short interfering RNA, useful for knock down of selected genes, by transforming cells with bank of short oligonucleotides and selection for expression of two reporter genes;  
     short oligonucleotide identification for short interfering RNA expression and specific gene knock down

L12 ANSWER 142 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI Identifying, analyzing and/or cloning nucleic acid isoforms, useful for preparing a probe, diagnosing a disease, or assessing responsiveness of a

patient to a treatment, comprises preparing complementary nucleic acid isoforms;

DNA isoform cloning and DNA probe for use in disease diagnosis

- |     |   |                  |              |
|-----|---|------------------|--------------|
| L12 | ANSWER 143 OF 314   | USPATFULL on STN | DUPLICATE 14 |
| TI  | Binary vectors for the improved transformation of plants systems  |                  |              |
| L12 | ANSWER 144 OF 314   | USPATFULL on STN | DUPLICATE 15 |
| TI  | Nucleic acid transfer vector for the introduction of nucleic acid into the DNA of a cell  |                  |              |
| L12 | ANSWER 145 OF 314   | USPATFULL on STN | DUPLICATE 16 |
| TI  | Bacteria with reduced genome  |                  |              |
| L12 | ANSWER 146 OF 314   | USPATFULL on STN | DUPLICATE 17 |
| TI  | Sequence specific recombinase-based methods for producing intron containing vectors and compositions for use in practicing the same |                  |              |
| L12 | ANSWER 147 OF 314   | USPATFULL on STN | DUPLICATE 18 |
| TI  | Site specific recombinase based method for producing adenoviral vectors   |                  |              |
| L12 | ANSWER 148 OF 314   | USPATFULL on STN | DUPLICATE 19 |
| TI  | Nuclease  |                  |              |
| L12 | ANSWER 149 OF 314   | USPATFULL on STN | DUPLICATE 20 |
| TI  | Recombinase-based methods for producing expression vectors and compositions for use in practicing the same                          |                  |              |
| L12 | ANSWER 150 OF 314   | USPATFULL on STN |              |
| TI  | Novel group B streptococcus antigens  |                  |              |
| L12 | ANSWER 151 OF 314   | USPATFULL on STN |              |
| TI  | Method for carrying out the parallel sequencing of a nucleic acid mixture on a surface  |                  |              |
| L12 | ANSWER 152 OF 314   | USPATFULL on STN |              |
| TI  | 207 human secreted proteins   |                  |              |
| L12 | ANSWER 153 OF 314   | USPATFULL on STN |              |
| TI  | Regulators of bacterial virulence factor expression   |                  |              |
| L12 | ANSWER 154 OF 314   | USPATFULL on STN |              |
| TI  | Anthrax synthase gene and method for increasing tryptophan production   |                  |              |
| L12 | ANSWER 155 OF 314   | USPATFULL on STN |              |
| TI  | Compositions and methods for the therapy and diagnosis of colon cancer  |                  |              |
| L12 | ANSWER 156 OF 314   | USPATFULL on STN |              |
| TI  | Novel oligonucleotide arrays and their use for sorting, isolating, sequencing, and manipulating nucleic acids                       |                  |              |
| L12 | ANSWER 157 OF 314   | USPATFULL on STN |              |
| TI  | Evolution of whole cells and organisms by recursive sequence recombination  |                  |              |
| L12 | ANSWER 158 OF 314   | USPATFULL on STN |              |
| TI  | Collections of binding proteins and tags and uses thereof for nested sorting and high throughput screening                          |                  |              |
| L12 | ANSWER 159 OF 314   | USPATFULL on STN |              |
| TI  | High throughput method for identification of sequence tags  |                  |              |

L12 ANSWER 160 OF 314 USPATFULL on STN  
 TI Attenuated human-bovine chimeric parainfluenza virus (PIV) vaccines

L12 ANSWER 161 OF 314 USPATFULL on STN  
 TI Tryparedoxin, expression plasmid, process of production, method of use, test kit, and pharmaceutical composition

L12 ANSWER 162 OF 314 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of pancreatic cancer

L12 ANSWER 163 OF 314 USPATFULL on STN  
 TI Human genes and gene expression products

L12 ANSWER 164 OF 314 USPATFULL on STN  
 TI NOVEL GROUP B STREPTOCOCCUS ANTIGENS

L12 ANSWER 165 OF 314 USPATFULL on STN  
 TI Cell type specific gene transfers using retroviral vectors containing antibody-envelope fusion proteins and wild-type envelope fusion proteins

L12 ANSWER 166 OF 314 USPATFULL on STN  
 TI Polypeptides and polynucleotides from coagulase-negative staphylococci

L12 ANSWER 167 OF 314 USPATFULL on STN  
 TI Nucleic acids encoding 3-ketoacyl-ACP reductase from Moraxella catarrhalis

L12 ANSWER 168 OF 314 USPATFULL on STN  
 TI Method for in vitro amplification of circular DNA

L12 ANSWER 169 OF 314 USPATFULL on STN  
 TI Assay for detecting apoptotic cells

L12 ANSWER 170 OF 314 USPATFULL on STN  
 TI Nucleic acid and amino acid sequences relating to Acinetobacter baumannii for diagnostics and therapeutics

L12 ANSWER 171 OF 314 USPATFULL on STN  
 TI Nucleotide sequence of the mycoplasma genitalium genome, fragments thereof, and uses thereof

L12 ANSWER 172 OF 314 USPATFULL on STN  
 TI Cell type specific gene transfer using retroviral vectors containing antibody-envelope fusion proteins and wild-type envelope fusion proteins

L12 ANSWER 173 OF 314 USPATFULL on STN  
 TI Nucleotide sequence of the Haemophilus influenzae Rd genome, fragments thereof, and uses thereof

L12 ANSWER 174 OF 314 USPATFULL on STN  
 TI Method of mapping restriction sites in polynucleotides

L12 ANSWER 175 OF 314 USPATFULL on STN  
 TI Nucleotide sequence of the Haemophilus influenzae Rd genome, fragments thereof, and uses thereof

L12 ANSWER 176 OF 314 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN  
 TI Distribution of Helicobacter pylori 51-specific genes on other Korean isolates of Helicobacter pylori.

L12 ANSWER 177 OF 314 USPATFULL on STN DUPLICATE 21  
 TI Anthranilate synthase gene and method of use thereof for conferring tryptophan overproduction

L12 ANSWER 178 OF 314 USPATFULL on STN DUPLICATE 22  
 TI Use of multiple recombination sites with unique specificity in recombinational cloning

L12 ANSWER 179 OF 314 USPATFULL on STN DUPLICATE 23  
 TI Soluble single chain T cell receptors

L12 ANSWER 180 OF 314 USPATFULL on STN  
 TI Bifunctional fusion proteins formed from hirudin and TAP

L12 ANSWER 181 OF 314 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of colon cancer

L12 ANSWER 182 OF 314 USPATFULL on STN  
 TI Fungal target genes and methods to identify those genes

L12 ANSWER 183 OF 314 USPATFULL on STN  
 TI Collections of binding proteins and tags and uses thereof for nested sorting and high throughput screening

L12 ANSWER 184 OF 314 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of ovarian cancer

L12 ANSWER 185 OF 314 USPATFULL on STN  
 TI Compositions and methods for the therapy and diagnosis of colon cancer

L12 ANSWER 186 OF 314 USPATFULL on STN  
 TI Tryparedoxin, expression plasmid, process of production, method of use, test kit, and pharmaceutical composition

L12 ANSWER 187 OF 314 USPATFULL on STN  
 TI Identification of congenital stationary night blindness in dogs

L12 ANSWER 188 OF 314 USPATFULL on STN  
 TI Recombinase-based methods for producing expression vectors and compositions for use in practicing the same

L12 ANSWER 189 OF 314 USPATFULL on STN  
 TI Computer readable genomic sequence of Haemophilus influenzae Rd, fragments thereof, and uses thereof

L12 ANSWER 190 OF 314 USPATFULL on STN  
 TI cDNAs coding for members of the carcinoembryonic antigen family

L12 ANSWER 191 OF 314 USP2 on STN  
 TI Methods for monitoring multiple gene expression

L12 ANSWER 192 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI New bacteriophage or plasmid cloning vectors, useful for in vitro or in vivo cloning nucleic acid inserts of interest used as tools in molecular genetic research;  
 vector-mediated reporter gene transfer and expression in host cell for gene analysis

L12 ANSWER 193 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI Determining a fraction of DNA molecules hemi-methylated at specific CpG dinucleotide sequence in a palindromic CpG methylation site, comprises

digesting DNA sample with an excess of methylation-sensitive restriction endonuclease;

DNA primer and polymerase chain reaction for methylation status evaluation

- L12 ANSWER 194 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
TI Novel oligonucleotide linker or population of linkers for preparing polynucleotide libraries, comprises an oligonucleotide fixed portion and an oligonucleotide variable portion;  
DNA primer and DNA sequencing for target ssDNA detection
- L12 ANSWER 195 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
TI Novel adaptor sequences for rapid joining with a target nucleic acid sequence, comprise topoisomerase recognition/cleavage sequence and a functional group or encoded functionality;  
DNA adaptor for polymerase chain reaction and target genome or DNA detection
- L12 ANSWER 196 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
TI Identification of the Staphylococcus aureus etd pathogenicity island which encodes a novel exfoliative toxin, ETD, and EDIN-B
- L12 ANSWER 197 OF 314 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN  
TI Type I restriction-modification systems from Lactobacillus delbrueckii subsp. lactis.
- L12 ANSWER 198 OF 314 USPATFULL on STN  
TI Oligonucleotide arrays and their use for sorting, isolating, sequencing, and manipulating nucleic acids
- L12 ANSWER 199 OF 314 USPATFULL on STN  
TI Pullulanase expression constructs containing  $\alpha$ -amylase promoter and leader sequences
- L12 ANSWER 200 OF 314 USPATFULL on STN  
TI Anthranilate synthase gene and method of use thereof for conferring tryptophan overproduction
- L12 ANSWER 201 OF 314 USPATFULL on STN  
TI Auxiliary genes and proteins of methicillin resistant bacteria and antagonists thereof
- L12 ANSWER 202 OF 314 USPATFULL on STN  
TI Method for introducing unidirectional nested deletions
- L12 ANSWER 203 OF 314 USPATFULL on STN  
TI Compositions, methods and kits for identifying naturally occurring RNA sequences having affinity for RNA-binding proteins
- L12 ANSWER 204 OF 314 USPATFULL on STN  
TI Compositions, methods, kits and apparatus for determining the presence or absence of target molecules
- L12 ANSWER 205 OF 314 USPATFULL on STN  
TI Identification of congenital stationary night blindness in dogs
- L12 ANSWER 206 OF 314 USPATFULL on STN  
TI Cystic fibrosis gene
- L12 ANSWER 207 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
TI Identification of a preferred liver transplant donor for donation to

patients with hepatitis C comprises determining the presence or absence of altered activity in a tumor necrosis factor;  
DNA primer and polymerase chain reaction for tumor necrosis factor polymorphism detection, genotyping and transplantation

- L12 ANSWER 208 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
TI DNA cleavage by type III restriction-modification enzyme EcoP15I is independent of spacer distance between two head to head oriented recognition sites
- L12 ANSWER 209 OF 314 USPATFULL on STN  
TI Auxiliary genes and proteins of methicillin resistant bacteria and antagonists thereof
- L12 ANSWER 210 OF 314 USPATFULL on STN  
TI Anthranilate synthase gene and method of use thereof for conferring tryptophan overproduction
- L12 ANSWER 211 OF 314 USPATFULL on STN  
TI Cleaved amplified modified polymorphic sequence detection methods
- L12 ANSWER 212 OF 314 USPATFULL on STN  
TI Method of sorting a mixture of nucleic acid strands on a binary array
- L12 ANSWER 213 OF 314 USPATFULL on STN  
TI Auxiliary genes and proteins of methicillin resistant bacteria and antagonists thereof
- L12 ANSWER 214 OF 314 USPATFULL on STN  
TI Compositions, methods, kits and apparatus for determining the presence or absence of protein component of telomerase enzyme
- L12 ANSWER 215 OF 314 USPATFULL on STN  
TI Plant promoter and method for gene expression using said promoter
- L12 ANSWER 216 OF 314 USPATFULL on STN  
TI cDNAs coding for members of the carcinoembryonic antigen family
- L12 ANSWER 217 OF 314 USPATFULL on STN  
TI Antibody preparations specifically binding to unique determinants of CEA antigens or fragments thereof and use of the antibody preparations in immunoassays
- L12 ANSWER 218 OF 314 USPATFULL on STN  
TI Auxiliary genes and proteins of methicillin resistant bacteria and antagonists thereof
- L12 ANSWER 219 OF 314 USPATFULL on STN  
TI Assay for detecting apoptotic cells
- L12 ANSWER 220 OF 314 LIFESCI COPYRIGHT 2009 CSA on STN DUPLICATE 24  
TI Broad-Range Bacteriophage Resistance in Streptococcus thermophilus by Insertional Mutagenesis
- L12 ANSWER 221 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 25  
TI Characterization of a Novel Plasmid-Encoded HsdS Subunit, S.LlaW12I, from Lactococcus lactis W12
- L12 ANSWER 222 OF 314 USPATFULL on STN  
TI Introns and exons of the cystic fibrosis gene and mutations thereof

L12 ANSWER 223 OF 314 USPATFULL on STN  
 TI Compositions, methods, kits and apparatus for determining the presence or absence of target molecules

L12 ANSWER 224 OF 314 USPATFULL on STN  
 TI Methods for screening for mutations at various positions in the introns and exons of the cystic fibrosis gene

L12 ANSWER 225 OF 314 USPATFULL on STN  
 TI Compound microsatellite primers for the detection of genetic polymorphisms

L12 ANSWER 226 OF 314 USPATFULL on STN  
 TI Fatty acid desaturase genes from plants

L12 ANSWER 227 OF 314 USPATFULL on STN  
 TI Cell type specific gene transfer using retroviral vectors containing antibody-envelope fusion proteins and wild-type envelope fusion proteins

L12 ANSWER 228 OF 314 USPATFULL on STN  
 TI Methods and materials for producing gene libraries

L12 ANSWER 229 OF 314 USPATFULL on STN  
 TI In vitro ligation of foreign DNA into large eukaryotic viruses

L12 ANSWER 230 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN  
 TI Phage resistance genes hsdR, hsdM and hsdS of lactic acid bacteria and recombinant bacteria producing the restriction endonuclease, methylase and HsdS

L12 ANSWER 231 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
 TI Regulation of endonuclease activity by proteolysis prevents breakage of unmodified bacterial chromosomes by type I restriction enzymes

L12 ANSWER 232 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN  
 TI DNA restriction dependent on two recognition sites: activities of the SfiI restriction-modification system in Escherichia coli

L12 ANSWER 233 OF 314 USPATFULL on STN  
 TI CDNA coding for carcinoembryonic antigen

L12 ANSWER 234 OF 314 USPATFULL on STN  
 TI Multimeric, recombinant urease vaccine

L12 ANSWER 235 OF 314 USPATFULL on STN  
 TI Method for producing lipolytic enzymes using transformed Pseudomonas

L12 ANSWER 236 OF 314 USPATFULL on STN  
 TI Polynucleotide sizing reagent

L12 ANSWER 237 OF 314 USPATFULL on STN  
 TI Methods of detecting cystic fibrosis gene by nucleic acid hybridization

L12 ANSWER 238 OF 314 USPATFULL on STN  
 TI Detection of nucleic acids in cells by thermophilic strand displacement amplification

L12 ANSWER 239 OF 314 USPATFULL on STN  
 TI Strand displacement amplification using thermophilic enzymes

L12 ANSWER 240 OF 314 USPATFULL on STN  
 TI Detection of nucleic acids in cells by thermophilic strand displacement

amplification

- L12 ANSWER 241 OF 314 USPATFULL on STN  
TI Isolated protein from *Eimeria* useful as a cross species vaccine
- L12 ANSWER 242 OF 314 USPATFULL on STN  
TI Nicking of DNA using boronated nucleotides
- L12 ANSWER 243 OF 314 USPATFULL on STN  
TI Method of inhibiting cell growth with the P.sub.20 receptor
- L12 ANSWER 244 OF 314 USPATFULL on STN  
TI Neuroblastoma-associated regulator gene
- L12 ANSWER 245 OF 314 USPATFULL on STN  
TI Methods and materials for producing gene libraries
- L12 ANSWER 246 OF 314 USPATFULL on STN  
TI Neuroblastoma-associated regulator gene
- L12 ANSWER 247 OF 314 USPATFULL on STN  
TI Strand displacement amplification using thermophilic enzymes
- L12 ANSWER 248 OF 314 USPATFULL on STN  
TI Detection of nucleic acids in cells by thermophilic strand displacement amplification
- L12 ANSWER 249 OF 314 USPATFULL on STN  
TI Methods of detecting compounds which bind to the P.sub.20 receptor
- L12 ANSWER 250 OF 314 USPATFULL on STN  
TI DNA Encoding the human P.sub.20 receptor and null cells expressing P.sub.20 receptors
- L12 ANSWER 251 OF 314 USPATFULL on STN  
TI Efficient directional genetic cloning system
- L12 ANSWER 252 OF 314 LIFESCI COPYRIGHT 2009 CSA on STN DUPLICATE 26  
TI Selection of non-specific DNA cleavage sites by the type IC restriction endonuclease EcoR124I
- L12 ANSWER 253 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 27  
TI The hsd loci of *Mycoplasma pulmonis*: organization, rearrangements and expression of genes
- L12 ANSWER 254 OF 314 USPATFULL on STN  
TI CDNA coding for carcinoembryonic antigen
- L12 ANSWER 255 OF 314 USPATFULL on STN  
TI Compositions and methods for making lipolytic enzymes
- L12 ANSWER 256 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
TI The methylation pattern of a cytosine DNA-methyltransferase gene *Arabidopsis thaliana* plants
- L12 ANSWER 257 OF 314 USPATFULL on STN  
TI Industrial yeast comprising an integrated glucoamylase gene
- L12 ANSWER 258 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN  
TI Highly efficient eukaryotic gene expression vectors for peptide secretion
- L12 ANSWER 259 OF 314 USPATFULL on STN



TI Expression and purification of recombinant soluble tissue factor  
 L12 ANSWER 260 OF 314 USPATFULL on STN  
 TI 25 KD coccidial antigen of elmeria tenella  
 L12 ANSWER 261 OF 314 USPATFULL on STN  
 TI Molecular cloning and expression of gene encoding lipolytic enzyme  
 L12 ANSWER 262 OF 314 USPATFULL on STN  
 TI cDNA coding for carcinoembryonic antigen (CEA)  
 L12 ANSWER 263 OF 314 USPATFULL on STN  
 TI Process and nucleic acid construct for producing reagent complexes useful in determining target nucleotide sequences  
 L12 ANSWER 264 OF 314 USPATFULL on STN  
 TI CDNAS coding for members of the carcinoembryonic antigen family  
 L12 ANSWER 265 OF 314 USPATFULL on STN  
 TI Recombinant baculovirus  
 L12 ANSWER 266 OF 314 USPATFULL on STN  
 TI DNA sequencing vector with reversible insert  
 L12 ANSWER 267 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 28  
 TI Macroevolution by transposition: Drastic modification of DNA recognition by a type I restriction enzyme following Tn5 transposition  
 L12 ANSWER 268 OF 314 USPATFULL on STN  
 TI CDNAS coding for members of the carcinoembryonic antigen family  
 L12 ANSWER 269 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN  
 TI Recombination of constant and variable modules alters DNA sequence recognition by type IC restriction - modification enzymes  
 L12 ANSWER 270 OF 314 USPATFULL on STN  
 TI Recombinant DNA molecules for producing terminal transferase-like polypeptides  
 L12 ANSWER 271 OF 314 USPATFULL on STN  
 TI Modified microorganisms and method of preparing and using same  
 L12 ANSWER 272 OF 314 USPATFULL on STN  
 TI Method and vector organism for controlled accumulation of cloned heterologous gene products in Bacillus subtilis  
 L12 ANSWER 273 OF 314 USPATFULL on STN  
 TI Yeast promoter and process for preparing heterologous protein  
 L12 ANSWER 274 OF 314 USPATFULL on STN  
 TI Hybrid interferons, their use as pharmaceutical compositions and as intermediate products for the preparation of antibodies and the use thereof and processes for preparing them  
 L12 ANSWER 275 OF 314 USPATFULL on STN  
 TI Novel expression control sequences  
 L12 ANSWER 276 OF 314 USPATFULL on STN  
 TI Method for cloning genes

L12 ANSWER 277 OF 314 USPATFULL on STN  
 TI Assay, reagent and kit employing nucleic acid strand displacement and restriction endonuclease cleavage

L12 ANSWER 278 OF 314 CAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 29  
 TI Evolution of DNA sequence specificity in type I restriction enzymes

L12 ANSWER 279 OF 314 USPATFULL on STN DUPLICATE 30  
 TI Methods and materials for obtaining microbial expression of polypeptides including bovine prolactin

L12 ANSWER 280 OF 314 USPATFULL on STN  
 TI Method and vector organism for controlled accumulation of cloned heterologous gene products in *Bacillus subtilis*

L12 ANSWER 281 OF 314 USPATFULL on STN  
 TI Recombinant DNA molecules and their use in producing human interferon-like polypeptides

L12 ANSWER 282 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
 TI EcoA and EcoE: Alternatives to the EcoK family of type I restriction and modification systems of *Escherichia coli*

L12 ANSWER 283 OF 314 LIFESCI COPYRIGHT 2009 CSA on STN DUPLICATE 31  
 TI Genetic recombination can generate altered restriction specificity.

L12 ANSWER 284 OF 314 USPATFULL on STN  
 TI Method for cloning genes

L12 ANSWER 285 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
 TI Structural homologies among type I restriction-modification system; an investigation to determine whether the systems are related in Enterobacteriaceae and *Escherichia coli* K12

L12 ANSWER 286 OF 314 USPATFULL on STN  
 TI Modified microorganisms and method of preparing and using same

L12 ANSWER 287 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
 TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof (PublishedApplication)

L12 ANSWER 288 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
 TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof (PublishedApplication)

L12 ANSWER 289 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
 TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof (PublishedApplication)

L12 ANSWER 290 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
 TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof (PublishedApplication)

L12 ANSWER 291 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
 TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof (PublishedApplication)

L12 ANSWER 292 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
 TI Methods for producing a paired tag from a nucleic acid sequence and methods of use thereof (PublishedApplication)



methods of use thereof (PublishedApplication)

L12 ANSWER 309 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and  
methods of use thereof (PublishedApplication)

L12 ANSWER 310 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and  
methods of use thereof (PublishedApplication)

L12 ANSWER 311 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and  
methods of use thereof (PublishedApplication)

L12 ANSWER 312 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and  
methods of use thereof (PublishedApplication)

L12 ANSWER 313 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and  
methods of use thereof (PublishedApplication)

L12 ANSWER 314 OF 314 USGENE COPYRIGHT 2009 SEQUENCEBASE CORP on STN  
TI Methods for producing a paired tag from a nucleic acid sequence and  
methods of use thereof (PublishedApplication)

-> d ibib abs 112 106 115 132 197 208 221 267 269 278 282 283 314

L12 ANSWER 106 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
ACCESSION NUMBER: 2005-29788 BIOTECHDS  
TITLE: New pure Type IIG restriction endonuclease obtainable from  
Citrobacter species or from Escherichia coli, useful for  
generating restriction endonucleases with new specificities;  
for use in genetic engineering  
AUTHOR: MORGAN R; WILSON G; LUNNEN K; HEITER D; BENNER J; NKENPOU C  
N; PICONE S  
PATENT ASSIGNEE: NEW ENGLAND BIOLABS INC  
PATENT INFO: WO 2005094516 13 Oct 2005  
APPLICATION INFO: WO 2005-US9824 23 Mar 2005  
PRIORITY INFO: US 2004-555796 24 Mar 2004; US 2004-555796 24 Mar 2004  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
OTHER SOURCE: WPI: 2005-714328 [73]  
AN 2005-29788 BIOTECHDS  
AB DERWENT ABSTRACT:

NOVELTY - A substantially pure Type IIG restriction  
endonuclease (I) obtainable from Citrobacter sp. 2144 (NEB#1398)  
(American Type Culture Collection (ATCC) Patent Accession Number PTA-5846)  
or from Escherichia coli NEB#1554 (ATCC Patent Accession Number PTA-5887),  
is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the  
following: (1) an isolated DNA obtainable from Citrobacter sp. 2144 (NEB  
(2) 1398) (ATCC Patent Accession Number PTA-5846) or from E.coli NEB (3)  
1554 (ATCC Patent Accession Number PTA-5887) and encoding (I), where the DNA  
comprises a first DNA segment expressing an endonuclease and methyl  
transferase catalytic function and a second DNA segment encoding a  
sequence specificity function of the restriction  
endonuclease, where the first and second DNA segments comprise  
one or more DNA molecules; (4) a recombinant DNA vector  
comprising at least one of first DNA segment coding for the  
restriction and modification domains of CspCI



psig. The lysate was centrifuged at 13000xG for 40 minutes and the supernatant collected. The supernatant solution was applied to a 400 ml diethylaminoethyl (DEAE) fast flow column. The diluted enzyme was applied to a 375 ml heparin hyper D column. A 2.5 L wash of buffer B was applied, then a 2 L gradient of sodium chloride from 0.15-1M in buffer B was applied and fractions were collected. Fractions were assayed for CspCI endonuclease activity by incubating with 1 microgram of phase lambda DNA (NEB) in 50 µl NEB buffer 2, supplemented with 20 micromolar for 15 minutes at 37degreesC. CspCI activity eluted at 0.3-0.35 M sodium chloride. CspCI activity eluted at 0.4-0.5 M potassium hydrogen phosphate. (87 pages)

L12 ANSWER 115 OF 314 USPATFULL on STN  
 ACCESSION NUMBER: 2004:280826 USPATFULL  
 TITLE: Methods and compositions relating to 5'-chimeric ribonucleic acids  
 INVENTOR(S): Sternberg, Paul, Pasadena, CA, UNITED STATES  
 Hwang, Byung Joon, San Marino, CA, UNITED STATES

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 20040220127	A1	20041104
APPLICATION INFO.:	US 2003-639016	A1	20030811 (10)

  

	NUMBER	DATE
PRIORITY INFORMATION:	US 2002-423490P	20021104 (60)
	US 2002-402473P	20020809 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	APPLICATION	
LEGAL REPRESENTATIVE:	ROPES & GRAY LLP, ONE INTERNATIONAL PLACE, BOSTON, MA, 02110-2624	
NUMBER OF CLAIMS:	57	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	20 Drawing Page(s)	
LINE COUNT:	3314	
CAS INDEXING IS AVAILABLE FOR THIS PATENT.		
AB The disclosure provides, among other things, methods for producing and using 5'-chimeric RNAs and cDNAs. 5'-chimeric RNAs and cDNAs may be used, for example, for high-throughput analysis of the 5'-end sequences for RNA transcripts.		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L12 ANSWER 132 OF 314 USPATFULL on STN  
 ACCESSION NUMBER: 2004:2072 USPATFULL  
 TITLE: Hybrid and single chain meganucleases and use thereof  
 INVENTOR(S): Arnould, Sylvain, Paris, FRANCE  
 Chames, Patrick, Paris, FRANCE  
 Choulika, Andre, Paris, FRANCE  
 Epinat, Jean-Charles, Paris, FRANCE  
 Lacroix, Emmanuel, Paris, FRANCE

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 20040002092	A1	20040101
APPLICATION INFO.:	US 2003-388230	A1	20030314 (10)

  

	NUMBER	DATE
PRIORITY INFORMATION:	US 2002-364113P	20020315 (60)
DOCUMENT TYPE:	Utility	

FILE SEGMENT: APPLICATION  
LEGAL REPRESENTATIVE: NIXON & VANDERHYE, PC, 1100 N GLEBE ROAD, 8TH FLOOR,  
ARLINGTON, VA, 22201-4714  
NUMBER OF CLAIMS: 26  
EXEMPLARY CLAIM: 1  
NUMBER OF DRAWINGS: 12 Drawing Page(s)  
LINE COUNT: 3746

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

AB This patent application relates to hybrid and/or single-chain rare-cutting endonucleases, called meganucleases, which recognize and cleave a specific nucleotide sequence, to polynucleotide sequences encoding for said rare-cutting endonucleases, to a vector comprising one of said polynucleotide sequences, to a cell or animal comprising one of said polynucleotide sequences or said rare-cutting endonucleases, to a process for producing one of said rare-cutting endonucleases and any use of the disclosed products and methods. More particularly, this invention contemplates any use of such rare-cutting endonuclease for genetic engineering and gene therapy.

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L12 ANSWER 197 OF 314 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on  
STN

ACCESSION NUMBER: 2002:585532 BIOSIS  
DOCUMENT NUMBER: PREV200200585532  
TITLE: Type I restriction-modification systems from *Lactobacillus delbrueckii* subsp. *lactis*.

AUTHOR(S): Bourniquel, A. A. [Reprint author]; Mollet, B.; Bickle, T. A. [Reprint author]

CORPORATE SOURCE: Biozentrum, Univ. of Basel, Basel, Switzerland  
SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (2002) Vol. 102, pp. 238. print.  
Meeting Info.: 102nd General Meeting of the American Society for Microbiology. Salt Lake City, UT, USA. May 19-23, 2002. American Society for Microbiology.  
ISSN: 1060-2011.

DOCUMENT TYPE: Conference; (Meeting)  
Conference; Abstract; (Meeting Abstract)

LANGUAGE: English

ENTRY DATE: Entered STN: 13 Nov 2002

Last Updated on STN: 13 Nov 2002

AB Background: *Lactobacillus delbrueckii* subsp. *lactis* is a lactic acid bacterium used worldwide for the production of Swiss-type hard cheeses. Whereas other dairy starters, e.g. *Lactococcus lactis* or *Streptococcus thermophilus*, are highly susceptible to bacteriophage infections, very few phages are known to target *L. delbrueckii* ssp. suggesting these bacteria possess a very active and reliable/versatile endogenous defense mechanism. Type I restriction-modification (R-M) systems are acknowledged defense mechanisms that depend on the generation of novel specificities for adaptability. Methods: Type I R-M hsd (host specificity for DNA) gene clusters were isolated from two strains of *L. delbrueckii* subsp. *lactis*, sequenced and characterized. In vivo transcription of the identified genes was verified by Northern blotting. The hsdR (restriction), hsdM (modification) and hsdS (specificity of DNA binding) genes were cloned into expression vectors for overexpression in *Escherichia coli*. The expressed proteins were purified, tested for activity and their recognition sites determined. Results: Both *L. delbrueckii* subsp. *lactis* strains examined possess type I R-M systems. The two hsd clusters (apprx8 kb) share a similar genetic organization consisting of: (i) the hsdR, hsdM and hsdS genes coding for a type I restriction enzyme, (ii) a second hsdS gene with a truncated 5'-end, and (iii) a gene similar to phage

integrases. The HsdR and HsdM subunits on both strains are highly conserved (98% identity), whereas the hsdS genes code for subunits with different specificities i.e. the two type I restriction enzymes have different recognition sites. Conclusion: Until recently, research on type I R-M systems focused on the enterobacteriaceae group in which hsd genes are but slightly conserved in-between strains. In the gram-positive bacterium *L. delbrueckii* subsp. *lactis* type I R-M genes are well-conserved, genetic variability being limited to the DNA binding domains of hsdS determining enzyme specificity. The presence of truncated hsdS and integrase genes in the hsd clusters suggests that novel specificities might be generated by domain shuffling.

L12 ANSWER 208 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
 ACCESSION NUMBER: 2001:33116733 BIOTECHNO  
 TITLE: DNA cleavage by type III restriction-modification enzyme EcoP15I is independent of spacer distance between two head to head oriented recognition sites  
 AUTHOR: Mucke M.; Reich S.; Moncke-Buchner E.; Reuter M.; Kruger D.H.  
 CORPORATE SOURCE: D.H. Kruger, Institut für Virologie, Medizinische Fakultät (Charité), Humboldt-Universität zu Berlin, D-10098, Berlin, Germany.  
 SOURCE: E-mail: detlev.kruger@charite.de  
 Journal of Molecular Biology, (28 SEP 2001), 312/4 (687-698), 42 reference(s)  
 CODEN: JMOBAK ISSN: 0022-2836  
 DOCUMENT TYPE: Journal; Article  
 COUNTRY: United Kingdom  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AN 2001:33116733 BIOTECHNO

AB The type III restriction-modification enzyme EcoP15I requires the interaction of two unmethylated, inversely oriented recognition sites 5'-CAG-CAG in head to head configuration to allow an efficient DNA cleavage. It has been hypothesized that two convergent DNA-translocating enzyme-substrate complexes interact to form the active cleavage complex and that translocation is driven by ATP hydrolysis. Using a half-automated, fluorescence-based detection method, we investigated how the distance between two inversely oriented recognition sites affects DNA cleavage efficiency. We determined that EcoP15I cleaves DNA efficiently even for two adjacent head to head or tail to tail oriented target sites. Hence, DNA translocation appears not to be required for initiating DNA cleavage in these cases. Furthermore, we report here that EcoP15I is able to cleave single-site substrates. When we analyzed the interaction of EcoP15I with DNA substrates containing adjacent target sites in the presence of non-hydrolyzable ATP analogues, we found that cleavage depended on the hydrolysis of ATP. Moreover, we show that cleavage occurs at only one of the two possible cleavage positions of an interacting pair of target sequences. When EcoP15I bound to a DNA substrate containing one recognition site in the absence of ATP, we observed a 36 nucleotide DNaseI-footprint that is asymmetric on both strands. All of our footprinting experiments showed that the enzyme did not cover the region around the cleavage site. Analyzing a DNA fragment with two head to head oriented recognition sites, EcoP15I protected 27-33 nucleotides around the recognition sequence, including an additional region of 26 bp between both cleavage sites. For all DNA substrates examined, the presence of ATP caused altered foot-printing patterns. We assume that the altered patterns are most likely due to a conformational change of the enzyme. Overall, our data further refine the tracking-collision model for type III restriction enzymes. .COPYRGHT. 2001 Academic Press.



COMMAND INTERRUPTED

If this message appears repeatedly, please notify the Help Desk.  
Enter "HELP STN" for information on contacting the nearest STN Help  
Desk by telephone or via SEND in the STNMAIL file.

-> d ibib abs 112 106 115 132 197 208 221 267 269 278 282 283 314

L12 ANSWER 106 OF 314 BIOTECHDS COPYRIGHT 2009 THOMSON REUTERS on STN  
ACCESSION NUMBER: 2005-29788 BIOTECHDS

TITLE: New pure Type IIG restriction endonuclease obtainable from  
Citrobacter species or from Escherichia coli, useful for  
generating restriction endonucleases with new specificities;  
for use in genetic engineering

AUTHOR: MORGAN R; WILSON G; LUNNEN K; HEITER D; BENNER J; NKENFOU C  
N; PICONE S

PATENT ASSIGNEE: NEW ENGLAND BIOLABS INC

PATENT INFO: WO 2005094516 13 Oct 2005

APPLICATION INFO: WO 2005-US9824 23 Mar 2005

PRIORITY INFO: US 2004-555796 24 Mar 2004; US 2004-555796 24 Mar 2004

DOCUMENT TYPE: Patent

LANGUAGE: English

OTHER SOURCE: WPI: 2005-714328 [73]

AN 2005-29788 BIOTECHDS

AB DERWENT ABSTRACT:

NOVELTY - A substantially pure Type IIG restriction  
endonuclease (I) obtainable from Citrobacter sp. 2144 (NEB#1398)  
(American Type Culture Collection (ATCC) Patent Accession Number PTA-5846)  
or from Escherichia coli NEB#1554 (ATCC Patent Accession Number PTA-5887),  
is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the  
following: (1) an isolated DNA obtainable from Citrobacter sp. 2144 (NEB  
(2) 1398) (ATCC Patent Accession Number PTA-5846) or from E.coli NEB (3)  
1554 (ATCC Patent Accession Number PTA-5887) and encoding (I), where the DNA  
comprises a first DNA segment expressing an endonuclease and methyl  
transferase catalytic function and a second DNA segment encoding a  
sequence specificity function of the restriction  
endonuclease, where the first and second DNA segments comprise  
one or more DNA molecules; (4) a recombinant DNA vector  
comprising at least one of first DNA segment coding for the  
restriction and modification domains of CspCI  
restriction endonuclease and a second segment coding  
for the specificity domain of the restriction  
endonuclease; (5) a host cell (II) transformed with a first DNA  
segment coding for the restriction and modification  
domains of CspCI restriction endonuclease and a  
second segment coding for the specificity domain of the  
restriction endonuclease, where the first DNA segment  
and the second DNA segment are contained with one or more DNA vectors;  
(6) preparing (I); and (7) making (MI) Type II restriction  
endonuclease having an altered specificity  
comprising: (a) selecting a restriction endonuclease  
from a set of enzymes, where each enzyme in the set is characterized by a  
modular structure having a specificity subunit and a catalytic subunit,  
the specificity subunit further comprising N-terminal domain for binding  
one half site of a bipartite recognition sequence and  
a C-terminal domain for binding a second half site of  
the bipartite recognition sequence; (b)  
modifying the specificity subunit; and (c) obtaining  
the Type II restriction endonuclease with  
altered specificity.

BIOTECHNOLOGY - Preparation: Preparing (I), involves cultivating a

sample of *Citrobacter* sp. 2144 (NEB#1398) or (II) under conditions favoring the production of the endonuclease, and purifying the endonuclease (claimed). Preferred Endonuclease: (I) Is capable of recognizing at least one sequence chosen (SEQ ID No: 32-35), and cleaving the DNA on both sides of the recognition sequence. Preferred Method: In (M1), modifying the specificity subunit further comprises: (a) substituting the N-terminal domain with a second C-terminal domain or substituting the C-terminal domain with a second N-terminal domain; (b) substituting the N-terminal domain or the C-terminal domain or both N-terminal and C-terminal domain with a binding domain from a second restriction endonuclease or methyltransferase; (c) mutating the N-terminal domain, the C-terminal domain or both domains to alter the binding specificity; or (d) changing the length of the spacer amino acid sequence between the N-terminal and C-terminal domains of the specificity module. The second restriction endonuclease or methyltransferase is chosen from Type I restriction endonuclease, Type IIG restriction endonuclease and gamma-type m6A methyltransferase. The specificity subunit and the catalytic subunit are encoded by different genes. (I) Comprises sequences such as  
 nnnnnnnnnncaannnnngtggnnnnnnnnnnnnnnnn (SEQ ID No: 32),  
 nnnnnnnnnncaannnnngtggnnnnnnnnnnnnnnnn (SEQ ID No: 33), caannnnnnngtgg (SEQ ID No: 34), caannnnngtgg (SEQ ID No: 35), where n=a, c, t or g.

USE - (I) Is useful for generating endonucleases with new specificity, for innovative genetic engineering.

EXAMPLE - CspCI was obtained by culturing either *Citrobacter* sp. 2144 (NEB#1398) or the transformed host *Escherichia coli* NEB#1554, and recovering the endonuclease from the cells. *Citrobacter* sp. 2144 (NEB#1398) or *E. coli* NEB#1554 were incubated aerobically at 37degreesC. Cells in the late logarithmic stage of growth were collected by centrifugation and either disrupted immediately or stored frozen at -70degreesC. The cell paste was suspended in a buffer solution and ruptured by sonication, high pressure dispersion or enzymatic digestion to allow extraction of the endonuclease by the buffer solution. Intact cells and cellular debris were then removed by centrifugation to produce a cell-free extract containing CspCI. The CspCI endonuclease was then purified from the cell-free extract by ion exchange chromatography, affinity chromatography, molecular sieve chromatography, or their combinations. 277 grams of *E. coli* NEB#1554 CspCI cell pellet or *Citrobacter* sp. 2144 were suspended in 1 liter of buffer A containing 300mM sodium chloride, and passed through a Gaulin homogenizer at 12000 psig. The lysate was centrifuged at 13000xG for 40 minutes and the supernatant collected. The supernatant solution was applied to a 400 ml diethylaminoethyl (DEAE) fast flow column. The diluted enzyme was applied to a 375 ml heparin hyper D column. A 2.5 L wash of buffer B was applied, then a 2 L gradient of sodium chloride from 0.15-1M in buffer B was applied and fractions were collected. Fractions were assayed for CspCI endonuclease activity by incubating with 1 microgram of phase lambda DNA (NEB) in 50 µl NEB buffer 2, supplemented with 20 micromolar for 15 minutes at 37degreesC. CspCI activity eluted at 0.3-0.35 M sodium chloride. CspCI activity eluted at 0.4-0.5 M potassium hydrogen phosphate. (87 pages)

L12 ANSWER 115 OF 314 USPATFULL on STN  
 ACCESSION NUMBER: 2004:280826 USPATFULL  
 TITLE: Methods and compositions relating to 5'-chimeric  
 ribonucleic acids  
 INVENTOR(S): Sternberg, Paul, Pasadena, CA, UNITED STATES  
 Hwang, Byung Joon, San Marino, CA, UNITED STATES

NUMBER	KIND	DATE
-----	-----	-----

PATENT INFORMATION: US 20040220127 A1 20041104  
 APPLICATION INFO.: US 2003-639016 A1 20030811 (10)

	NUMBER	DATE
PRIORITY INFORMATION:	US 2002-423490P	20021104 (60)
	US 2002-402473P	20020809 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	APPLICATION	
LEGAL REPRESENTATIVE:	ROPES & GRAY LLP, ONE INTERNATIONAL PLACE, BOSTON, MA, 02110-2624	
NUMBER OF CLAIMS:	57	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	20 Drawing Page(s)	
LINE COUNT:	3314	
CAS INDEXING IS AVAILABLE FOR THIS PATENT.		
AB The disclosure provides, among other things, methods for producing and using 5'-chimeric RNAs and cDNAs. 5'-chimeric RNAs and cDNAs may be used, for example, for high-throughput analysis of the 5'-end sequences for RNA transcripts.		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L12 ANSWER 132 OF 314 USPTAFULL on STN  
 ACCESSION NUMBER: 2004:2072 USPTAFULL  
 TITLE: Hybrid and single chain meganucleases and use thereof  
 INVENTOR(S): Arnould, Sylvain, Paris, FRANCE  
 Chames, Patrick, Paris, FRANCE  
 Choulaka, Andre, Paris, FRANCE  
 Epinat, Jean-Charles, Paris, FRANCE  
 Lacroix, Emmanuel, Paris, FRANCE

	NUMBER	KIND	DATE
PATENT INFORMATION:	US 20040002092	A1	20040101
APPLICATION INFO.:	US 2003-388230	A1	20030314 (10)

	NUMBER	DATE
PRIORITY INFORMATION:	US 2002-364113P	20020315 (60)
DOCUMENT TYPE:	Utility	
FILE SEGMENT:	APPLICATION	
LEGAL REPRESENTATIVE:	NIXON & VANDERHYE, PC, 1100 N GLEBE ROAD, 8TH FLOOR, ARLINGTON, VA, 22201-4714	
NUMBER OF CLAIMS:	26	
EXEMPLARY CLAIM:	1	
NUMBER OF DRAWINGS:	12 Drawing Page(s)	
LINE COUNT:	3746	
CAS INDEXING IS AVAILABLE FOR THIS PATENT.		
AB This patent application relates to hybrid and/or single-chain rare-cutting endonucleases, called meganucleases, which recognize and cleave a specific nucleotide sequence, to polynucleotide sequences encoding for said rare-cutting endonucleases, to a vector comprising one of said polynucleotide sequences, to a cell or animal comprising one of said polynucleotide sequences or said rare-cutting endonucleases, to a process for producing one of said rare-cutting endonucleases and any use of the disclosed products and methods. More particularly, this invention contemplates any use of such rare-cutting endonuclease for genetic engineering and gene therapy.		

CAS INDEXING IS AVAILABLE FOR THIS PATENT.

L12 ANSWER 197 OF 314 BIOSIS COPYRIGHT (c) 2009 The Thomson Corporation on STN

ACCESSION NUMBER: 2002:585532 BIOSIS  
DOCUMENT NUMBER: PREV200200585532  
TITLE: Type I restriction-modification systems from *Lactobacillus delbrueckii* subsp. *lactis*.  
AUTHOR(S): Bourniquel, A. A. [Reprint author]; Mollet, B.; Bickle, T. A. [Reprint author]  
CORPORATE SOURCE: Biozentrum, Univ. of Basel, Basel, Switzerland  
SOURCE: Abstracts of the General Meeting of the American Society for Microbiology, (2002) Vol. 102, pp. 238. print.  
Meeting Info.: 102nd General Meeting of the American Society for Microbiology. Salt Lake City, UT, USA. May 19-23, 2002. American Society for Microbiology.  
ISSN: 1060-2011.  
DOCUMENT TYPE: Conference; (Meeting)  
Conference; Abstract; (Meeting Abstract)  
LANGUAGE: English  
ENTRY DATE: Entered STN: 13 Nov 2002  
Last Updated on STN: 13 Nov 2002

AB Background: *Lactobacillus delbrueckii* subsp. *lactis* is a lactic acid bacterium used worldwide for the production of Swiss-type hard cheeses. Whereas other dairy starters, e.g. *Lactococcus lactis* or *Streptococcus thermophilus*, are highly susceptible to bacteriophage infections, very few phages are known to target *L. delbrueckii* ssp. suggesting these bacteria possess a very active and reliable/versatile endogenous defense mechanism. Type I restriction-modification (R-M) systems are acknowledged defense mechanisms that depend on the generation of novel specificities for adaptability. Methods: Type I R-M hsd (host specificity for DNA) gene clusters were isolated from two strains of *L. delbrueckii* subsp. *lactis*, sequenced and characterized. In vivo transcription of the identified genes was verified by Northern blotting. The hsdR (restriction), hsdM (modification) and hsdS (specificity of DNA binding) genes were cloned into expression vectors for overexpression in *Escherichia coli*. The expressed proteins were purified, tested for activity and their recognition sites determined. Results: Both *L. delbrueckii* subsp. *lactis* strains examined possess type I R-M systems. The two hsd clusters (apprx 8 kb) share a similar genetic organization consisting of: (i) the hsdR, hsdM and hsdS genes coding for a type I restriction enzyme, (ii) a second hsdS gene with a truncated 5'-end, and (iii) a gene similar to phage integrases. The HsdR and HsdM subunits on both strains are highly conserved (98% identity), whereas the hsdS genes code for subunits with different specificities i.e. the two type I restriction enzymes have different recognition sites. Conclusion: Until recently, research on type I R-M systems focused on the enterobacteriaceae group in which hsd genes are but slightly conserved in-between strains. In the gram-positive bacterium *L. delbrueckii* subsp. *lactis* type I R-M genes are well-conserved, genetic variability being limited to the DNA binding domains of hsdS determining enzyme specificity. The presence of truncated hsdS and integrases genes in the hsd clusters suggests that novel specificities might be generated by domain shuffling.

L12 ANSWER 208 OF 314 BIOTECHNO COPYRIGHT 2009 Elsevier Science B.V. on STN  
ACCESSION NUMBER: 2001:33116733 BIOTECHNO

TITLE: DNA cleavage by type III restriction-modification enzyme EcoP15I is independent of spacer distance between two head to head oriented recognition sites  
AUTHOR: Mucke M.; Reich S.; Moncke-Buchner E.; Reuter M.; Kruger D.H.  
CORPORATE SOURCE: D.H. Kruger, Institut für Virologie, Medizinische Fakultät (Charité), Humboldt-Universität zu Berlin,

D-10098, Berlin, Germany.  
 E-mail: detlev.kruger@charite.de  
 SOURCE: Journal of Molecular Biology, (28 SEP 2001), 312/4  
 (687-698), 42 reference(s)  
 CODEN: JMOBAK ISSN: 0022-2836  
 DOCUMENT TYPE: Journal; Article  
 COUNTRY: United Kingdom  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English  
 AN 2001:33116733 BIOTECHNO  
 AB The type III restriction-modification enzyme

AB The type III restriction-modification enzyme EcoP15I requires the interaction of two unmethylated, inversely oriented recognition sites 5'-CAG-CAG in head to head configuration to allow an efficient DNA cleavage. It has been hypothesized that two convergent DNA-translocating enzyme-substrate complexes interact to form the active cleavage complex and that translocation is driven by ATP hydrolysis. Using a half-automated, fluorescence-based detection method, we investigated how the distance between two inversely oriented recognition sites affects DNA cleavage efficiency. We determined that EcoP15I cleaves DNA efficiently even for two adjacent head to head or tail to tail oriented target sites. Hence, DNA translocation appears not to be required for initiating DNA cleavage in these cases. Furthermore, we report here that EcoP15I is able to cleave single-site substrates. When we analyzed the interaction of EcoP15I with DNA substrates containing adjacent target sites in the presence of non-hydrolyzable ATP analogues, we found that cleavage depended on the hydrolysis of ATP. Moreover, we show that cleavage occurs at only one of the two possible cleavage positions of an interacting pair of target sequences. When EcoP15I bound to a DNA substrate containing one recognition site in the absence of ATP, we observed a 36 nucleotide *DnaS*I-footprint that is asymmetric on both strands. All of our footprinting experiments showed that the enzyme did not cover the region around the cleavage site. Analyzing a DNA fragment with two head to head oriented recognition sites, EcoP15I protected 27-33 nucleotides around the recognition sequence, including an additional region of 26 bp between both cleavage sites. For all DNA substrates examined, the presence of ATP caused altered foot-printing patterns. We assume that the altered patterns are most likely due to a conformational change of the enzyme. Overall, our data further refine the tracking-collision model for type III restriction enzymes. .COPYRG. 2001 Academic Press.

COMMAND INTERRUPTED

```
=> d ibib abs 112 208 221 267 269 278 282 283 314
```

DOCUMENT TYPE: Journal; Article  
COUNTRY: United Kingdom  
LANGUAGE: English  
SUMMARY LANGUAGE: English  
AN 2001:33116733 BIOTECHNO

AB The type III restriction-modification enzyme EcoP15I requires the interaction of two unmethylated, inversely oriented recognition sites 5'-CAG-CAG in head to head configuration to allow an efficient DNA cleavage. It has been hypothesized that two convergent DNA-translocating enzyme-substrate complexes interact to form the active cleavage complex and that translocation is driven by ATP hydrolysis. Using a half-automated, fluorescence-based detection method, we investigated how the distance between two inversely oriented recognition sites affects DNA cleavage efficiency. We determined that EcoP15I cleaves DNA efficiently even for two adjacent head to head or tail to tail oriented target sites. Hence, DNA translocation appears not to be required for initiating DNA cleavage in these cases. Furthermore, we report here that EcoP15I is able to cleave single-site substrates. When we analyzed the interaction of EcoP15I with DNA substrates containing adjacent target sites in the presence of non-hydrolyzable ATP analogues, we found that cleavage depended on the hydrolysis of ATP. Moreover, we show that cleavage occurs at only one of the two possible cleavage positions of an interacting pair of target sequences. When EcoP15I bound to a DNA substrate containing one recognition site in the absence of ATP, we observed a 36 nucleotide DNaseI-footprint that is asymmetric on both strands. All of our footprinting experiments showed that the enzyme did not cover the region around the cleavage site. Analyzing a DNA fragment with two head to head oriented recognition sites, EcoP15I protected 27-33 nucleotides around the recognition sequence, including an additional region of 26 bp between both cleavage sites. For all DNA substrates examined, the presence of ATP caused altered foot-printing patterns. We assume that the altered patterns are most likely due to a conformational change of the enzyme. Overall, our data further refine the tracking-collision model for type III restriction enzymes. .COPYRG. 2001 Academic Press.

COMMAND INTERRUPTED

If this message appears repeatedly, please notify the Help Desk.  
Enter "HELP STN" for information on contacting the nearest STN Help Desk by telephone or via SEND in the STNMAIL file.

-> d ibib abs 112 221 267 269 278 282 283 314

COMMAND INTERRUPTED

If this message appears repeatedly, please notify the Help Desk.  
Enter "HELP STN" for information on contacting the nearest STN Help Desk by telephone or via SEND in the STNMAIL file.

-> d ibib abs 112 267 269 278 282 283 314

COMMAND INTERRUPTED

If this message appears repeatedly, please notify the Help Desk.  
Enter "HELP STN" for information on contacting the nearest STN Help Desk by telephone or via SEND in the STNMAIL file.

=> d his full

(FILE 'HOME' ENTERED AT 00:49:28 ON 01 FEB 2009)

INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, ANTE, AQUALINE,

AQUASCI, BIOENG, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CAPLUS,  
 CEABA-VTB, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE, DISSABS, DRUGB,  
 DRUGMONO2, DRUG, EMBAL, EMBASE, ...' ENTERED AT 00:50:04 ON 01 FEB 2009  
 SEA (RESTRIC? (3A)ENDONUCLEAS?) OR (RESTRIC? (3A)ENZYM?) OR (REST

```

-----
21  FILE ADISCTI
3   FILE ADISINSIGHT
7   FILE ADISNEWS
3845 FILE AGRICOLA
104  FILE ANABSTR
42   FILE ANTE
40   FILE AQUALINE
1189 FILE AQUASCI
3479 FILE BIOENG
31117 FILE BIOSIS
10212 FILE BIOTECHABS
10212 FILE BIOTECHDS
17093 FILE BIOTECHNO
10248 FILE CABA
40802 FILE CAPLUS
731  FILE CEABA-VTB
77   FILE CIN
338  FILE CONFSCI
3    FILE CROPB
124  FILE CROPU
19   FILE DDFB
133  FILE DDFU
43989 FILE DGENE
2088 FILE DISSABS
19   FILE DRUGB
425  FILE DRUGU
88   FILE EMBAL
21832 FILE EMBASE
9157 FILE ESBIOBASE
363  FILE FROSTI
1154 FILE FSTA
2282317 FILE GENBANK
53   FILE HEALSFE
7069 FILE IPIPAT
11   FILE IMSDRUGNEWS
9    FILE IMSRESEARCH
20   FILE KOSMET
17321 FILE LIPESCI
40206 FILE MEDLINE
199  FILE NTIS
348  FILE OCEAN
11890 FILE PASCAL
164  FILE PCTGEN
1    FILE PHAR
1    FILE PHARMAML
66   FILE PHIN
640  FILE PROMT
1    FILE PROUSDDR
3    FILE RDISCLOSURE
20911 FILE SCISEARCH
10529 FILE TOXCENTER
14486 FILE USGENE
72194 FILE USPATFULL
29   FILE USPATOLD
11872 FILE USPAT2
1    FILE VETB
144  FILE VETU

```

```

        62   FILE WATER
    9109   FILE WPIDS
        62   FILE WPIFV
    9109   FILE WPINDEX
        37   FILE IPA
        4    FILE NAPRALERT
    488    FILE NLDB
L1        QUE (RESTRIC?(3A) ENDONUCLEAS?) OR (RESTRIC?(3A) ENZYM?) OR
        (RESTRIC?(3A) MODIF?(5A) (ENZYM? OR ENDONUCLEAS? OR SYSTEM?))
-----
        D RANK

FILE 'USPATFULL, CAPLUS, MEDLINE, BIOSIS, EMBASE, SCISEARCH, LIFESCI,
BIOTECHNO, USGENE, PASCAL, USPAT2, TOXCENTER, CABA, BIOTECHDS' ENTERED AT
01:00:54 ON 01 FEB 2009
L2        330713 SEA (RESTRIC?(3A) ENDONUCLEAS?) OR (RESTRIC?(3A) ENZYM?) OR
        (RESTRIC?(3A) MODIF?(5A) (ENZYM? OR ENDONUCLEAS? OR SYSTEM?))
L3        79162 SEA L2(S) (SPECIFIC? OR RECOG?(S) (SEQUENC? OR DNA?))
L4        49041 SEA L3 AND (HYBRID? OR RECOMBINAT? OR TRUNCAT? OR TRANSPOS?)
L5        990 SEA L3(S) ((TWO(3A) RECOGNIT?(3A) SITE?) OR HSDS?)
L6        344 SEA L5 (S) (HYBRID? OR RECOMBIN? OR TRUNCAT? OR EXCHANG? OR
        TRANSPOS? OR ALTER?)
L7        280 DUP REM L6 (64 DUPLICATES REMOVED)
L8        14 SEA L7(S) HALF?
        D TI L8 1-14
        D L8 IBIB ABS 8 12 13
L9        1657 SEA L2(S) ((TWO(3A) RECOG?(3A) SITE?) OR HSDS? OR (HALF?(3A)
        SITE?))
L10       500 SEA L9(S) (HYBRID? OR RECOMB? OR TRUNC? OR EXCH? OR TRANSPO? OR
        ALTER?)
L11       359 SEA L10 AND ((MODIF? OR ALTER? OR HYBRID?) (3A) (DNA? OR SEQUEN?
        OR SPECIFIC? OR (RECOGN?(3A) SITE?)))
L12       314 DUP REM L11 (45 DUPLICATES REMOVED)
        D TI L12 1-314
        D IBIB ABS L12 106 115 132 197 208 221 267 269 278 282 283 314
        D IBIB ABS L12 106 115 132 197 208 221 267 269 278 282 283 314
        D IBIB ABS L12 208 221 267 269 278 282 283 314
        D IBIB ABS L12 221 267 269 278 282 283 314
        D IBIB ABS L12 267 269 278 282 283 314

```

FILE HOME

FILE STINDEX

FILE USPATFULL

FILE COVERS 1971 TO PATENT PUBLICATION DATE: 29 Jan 2009 (20090129/PD)

FILE LAST UPDATED: 29 Jan 2009 (20090129/ED)

HIGHEST GRANTED PATENT NUMBER: US7484247

HIGHEST APPLICATION PUBLICATION NUMBER: US20090031463

CA INDEXING IS CURRENT THROUGH 29 Jan 2009 (20090129/UPCA)

ISSUE CLASS FIELDS (/INCL) CURRENT THROUGH: 29 Jan 2009 (20090129/PD)

REVISED CLASS FIELDS (/NCL) LAST RELOADED: Dec 2008

USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Dec 2008

USPATFULL now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

FILE CAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available



for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 1 Feb 2009 VOL 150 ISS 6  
FILE LAST UPDATED: 29 Jan 2009 (20090129/ED)

Capius now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

FILE MEDLINE  
FILE LAST UPDATED: 31 Jan 2009 (20090131/UP). FILE COVERS 1949 TO DATE.

MEDLINE and IMEDLINE have been updated with the 2009 Medical Subject Headings (MeSH) vocabulary and tree numbers from the U.S. National Library of Medicine (NLM). Additional information is available at

[http://www.nlm.nih.gov/pubs/techbull/nd08/nd08\\_medline\\_data\\_changes\\_2009](http://www.nlm.nih.gov/pubs/techbull/nd08/nd08_medline_data_changes_2009).

This file contains CAS Registry Numbers for easy and accurate substance identification.

See HELP RANGE before carrying out any RANGE search.

MEDLINE Accession Numbers (ANs) for records from 1950-1977 have been converted from 8 to 10 digits. Searches using an 8 or 10 digit AN will retrieve the same record. The 10-digit ANs can be expanded, searched, and displayed in all records from 1949 to the present.

FILE BIOSIS  
FILE COVERS 1926 TO DATE.  
CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT  
FROM JANUARY 1926 TO DATE.

RECORDS LAST ADDED: 28 January 2009 (20090128/ED)

BIOSIS has been augmented with 1.8 million archival records from 1926 through 1968. These records have been re-indexed to match current BIOSIS indexing.

FILE EMBASE  
FILE COVERS 1974 TO 30 Jan 2009 (20090130/ED)

EMBASE was reloaded on March 30, 2008.

EMBASE is now updated daily. SDI frequency remains weekly (default) and biweekly.

This file contains CAS Registry Numbers for easy and accurate substance identification.

Beginning January 2008, Elsevier will no longer provide EMTREE codes as part of the EMTREE thesaurus in EMBASE. Please update your current-awareness alerts (SDIs) if they contain EMTREE codes.

For further assistance, please contact your local helpdesk.

FILE SCISEARCH

FILE COVERS 1974 TO 29 Jan 2009 (20090129/ED)

SCISEARCH has been reloaded, see HELP RLOAD for details.

FILE LIFESCI

FILE COVERS 1978 TO 20 Jan 2009 (20090120/ED)

FILE BIOTECHNO

FILE LAST UPDATED: 7 JAN 2004 <20040107/UP>

FILE COVERS 1980 TO 2003.

THIS FILE IS A STATIC FILE WITH NO UPDATES

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION AVAILABLE IN  
/CT AND BASIC INDEX <<<

FILE USGENE

FILE LAST UPDATED: 30 JAN 2009 <20090130/UP>

MOST RECENT PUBLICATION DATE: 15 JAN 2009 <20090115/PD>

FILE COVERS 1982 TO DATE

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION (SLRT) IS AVAILABLE  
IN THE BASIC INDEX (/BI) AND FEATURE TABLE (/FEAT) FIELDS <<<

>>> DOWNLOAD THE USGENE WORKSHOP MANUAL:

[http://www.stn-international.com/USGENE\\_workshop\\_manual.html](http://www.stn-international.com/USGENE_workshop_manual.html)

>>> DOWNLOAD RUN BLAST/GETSIM FREQUENTLY ASKED QUESTIONS:

<http://www.stn-international.com/usgenefaq.html> <<<

>>> DOWNLOAD COMPLETE USGENE HELP AS PDF:

[http://www.stn-international.com/usgene\\_help.html](http://www.stn-international.com/usgene_help.html) <<<

>>> USGENE now provides USPTO sequence data within 3 days of publication  
- see NEWS <<<

>>> SEARCH AND DISPLAY OF USPTO EXEMPLARY CLAIM (ECLM) IS AVAILABLE !! <<<

>>> NEW SEQUENCE SEARCH INTERACTION TO REFINE ANSWER

SETS BY PERCENT (%) NOW AVAILABLE - TO LEARN MORE, VISIT:

[http://www.stn-international.com/New\\_sequence\\_search.html](http://www.stn-international.com/New_sequence_search.html) <<<

FILE PASCAL

FILE LAST UPDATED: 26 JAN 2009 <20090126/UP>

FILE COVERS 1977 TO DATE.

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION IS AVAILABLE  
IN THE BASIC INDEX (/BI) FIELD <<<

FILE USPAT2

FILE COVERS 2001 TO PUBLICATION DATE: 29 Jan 2009 (20090129/PD)

FILE LAST UPDATED: 29 Jan 2009 (20090129/ED)

HIGHEST GRANTED PATENT NUMBER: US20080088069  
HIGHEST APPLICATION PUBLICATION NUMBER: US20090030948  
CA INDEXING IS CURRENT THROUGH 29 Jan 2009 (20090129/UPCA)  
ISSUE CLASS FIELDS (/INCL) CURRENT THROUGH: 29 Jan 2009 (20090129/PD)  
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Dec 2008  
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Dec 2008

USPAT2 now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

#### FILE TOXCENTER

FILE COVERS 1907 TO 27 Jan 2009 (20090127/ED)

The MEDLINE file segment has been updated with the National Library of Medicine's revised 2008 MeSH terms. See HELP RLOAD for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

The BIOSIS segment of TOXCENTER has been augmented with 13,000 records from 1946 through 1968.

#### FILE CABA

FILE COVERS 1973 TO 8 Jan 2009 (20090108/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

The CABA file was reloaded 7 December 2003. Enter HELP RLOAD for details.

#### FILE BIOTECHDS

FILE LAST UPDATED: 12 JAN 2009 <20090112/UP>

FILE COVERS 1982 TO DATE

>>> USE OF THIS FILE IS LIMITED TO BIOTECH SUBSCRIBERS <<<

=> d ibib abs 112 283

L12 ANSWER 283 OF 314 LIFESCI COPYRIGHT 2009 CSA on STN DUPLICATE 31

ACCESSION NUMBER: 84:97138 LIFESCI

TITLE: Genetic recombination can generate altered restriction specificity.

AUTHOR: Fuller-Pace, F.V.; Bullas, L.R.; Delius, H.; Murray, N.E.  
CORPORATE SOURCE: Dep. Mol. Biol., Univ. Edinburgh, King's Build., Mayfield Rd., Edinburgh EH9 3JR, UK

SOURCE: PROC. NATL. ACAD. SCI. USA., (1984) vol. 81, no. 19, pp. 6095-6099.

DOCUMENT TYPE: Journal

FILE SEGMENT: N; G

LANGUAGE: English

SUMMARY LANGUAGE: English

AB A recombinant strain, isolated following the transduction of an *Escherichia coli* recipient carrying the *Salmonella typhimurium* (SB) specificity genes with DNA from a donor having the *Salmonella potsdam* (SP) specificity, was shown to have neither SB nor SP specificity but to encode a novel restriction specificity, SQ. The heteroduplex analysis of the *hsdS* (specificity) genes of the SB and SP restriction and modification systems described here identifies a conserved sequence of around 100 base pairs flanked by two nonhomologous regions each of approximately 500 base pairs. The